



**MODULE - 1** Diversity and Evolution of Life



## TISSUES AND OTHER LEVELS OF ORGANIZATION

You have just learnt that cell is the fundamental structural and functional unit of organisms and that bodies of organisms are made up of cells of various shapes and sizes. Groups of similar cells aggregate to collectively perform a particular function. Such groups of cells are termed "tissues". This lesson deals with the various kinds of tissues of plants and animals.



After completing this lesson, you will be able to :

- *define tissues;*
- classify plant tissues;
- name the various kinds of plant tissues;
- enunciate the tunica corpus theory and histogen theory;
- classify animal tissues;
- describe the structure and function of various kinds of epithelial tissues;
- describe the structure and function of various kinds of connective tissues;
- describe the structure and function of muscular tissue;
- describe the structure and function of nervous tissue.

#### 5.1 WHAT IS A TISSUE

Organs such as stem, and roots in plants, and stomach, heart and lungs in animals are made up of different kinds of tissues. A **tissue is a group of cells with a common origin, structure and function.** Their common origin means they are derived from the same layer (details in lesson No. 20) of cells in the embryo. Being of a common origin, there are similar in structure and hence perform the same function. Several types of tissues organise to form an **organ**.

**Example :** Blood, bone, and cartilage are some examples of animal tissues whereas parenchyma, collenchyma, xylem and phloem are different tissues present in the plants. The study of tissues is called **histology**.

### **MODULE - 1** Diversity and Evolution

of Life



A group of cells with similar origin, structure and function is called **tissue**. e.g. bone, and muscle in animals and meristem in tips of root and shoot in plants

#### **5.2 THE PLANT TISSUES**

The plant tissues are mainly of two catagories:

- 1. Meristematic (Gk. meristos : dividing)
- 2. Permanent (non-dividing)

#### 1. Meristematic tissues

- Composed of immature or undifferentiated cells without intercellular spaces.
- The cells may be rounded, oval or polygonal; always living and thin-walled.
- Each cell has abundant cytoplasm and a prominent nucleus in it.
- Vacuoles may be small or absent.

#### Table 5.1 Types of meristematic tissue

Types	Location	Function
Apical Meristem	Root tip and shoot tip.	Growth in length of plants and their branches.
Intercalary Meristem	At the bases of leaves or at the bases of internodes.	Internodal growth, in monocots growth of leaf lamina in grasses.
Lateral Meristem	Cambium between xylem and phloem and cork. cambium in the cortex of dicot plants.	Growth in thickness of the plant body (secondary growth).



#### 2. Permanent tissues

- Permanent tissues are those in which growth has stopped either completely or for the time being.
- Cells of these tissues may be living or dead; and thin-walled or thickwalled.
- Thin-walled permanent tissues are generally living whereas the thick-walled tissues may be living or dead.

#### Types of permanent tissues

- (i) **Simple tissues :** A simple tissue is made up of only one type of cells. Common simple tissues are parenchyma, collenchyma and sclerenchyma (Fig. 5.2, 5.3 and 5.4).
- (ii) **Complex tissues :** A complex tissue is made up of more than one type of cells working together as a unit. Common examples are xylem and phloem (Fig. 5.5 and 5.6).

The structure, function and distribution of simple plant tissues is given in Table 5.2.



1. Define a tissue.

2. Give one word equivalent for the following :

(i) A plant tissue that consists of cells which continue to divide to produce more cells.

- .....
- (ii) The meristematic tissue responsible for the increase in thickness of the stem of a tree.

.....

(iii) The kind of plant tissues which consists of all similar cells.

.....

.....

- (iv) The category of plant tissues in which the cells do not divide.
- 3. What do you mean by "cells of a tissue have similar origin"?
- -----
- 4. Name that branch of Biology in which tissues are studied?
  - -----
- 5. What is a complex tissue?
- -----

6. Mention any **two** special features of the meristematic cells.

**BIOLOGY** 



**Notes** 

MODULE - 1

Diversity and Evolution of Life

#### **MODULE - 1** Diversity and Evolution

of Life

5.2.1 Simple Plant Tissues

There are three types of simple plant tissues (Fig. 5.2, 5.3 and 5.4)

- 1. Parenchyma (Chlorenchyma and Aerenchyma)
- 2. Collenchyma
- 3. Sclerenchyma



#### Table 5.2 Structure, Function and Distribution of simple tissues

Tissue	Living or Dead	Structure	Function	Distribution
1. Parenchyma	Living	<ul> <li>(i) Oval or round, thin-walled with sufficient cytoplasm.</li> <li>(ii) Has prominent nucleus and intercellular spaces</li> <li>(iii) Wall made up of cellulose</li> </ul>	<ul> <li>(a) They make large parts of various organs in most plants.</li> <li>(b) Act as storage cells.</li> <li>(c) Chlorenchyma carries out photosynthesis.</li> </ul>	<ol> <li>Pith and cortex of stem and root.</li> <li>Mesophyll of leaves.</li> <li>Endosperm of seed.</li> <li>Xylem and phloam</li> </ol>
<ul><li>(a) Chloren- chyma</li><li>(b) Aeren- chyma</li></ul>	Living Living	Parenchyma containing chloroplasts. Parenchyma with large air spaces or	(d) Turgid, parenchyma gives rigidity to the plant body.	parenchyma in vascular tissue. 5. Occur in
2. Collenchyma (Gk. collen : glue)	Living	<ul> <li>intercellular spaces.</li> <li>(i) Elongated cells with thick primary walls. Thickenings more in the corners of the cells.</li> <li>(ii) Wall material is cellulose and pectin</li> <li>(iii) Intercellular sapces present.</li> </ul>	Gives mechanical support to the plant body. Specially in many dicot leaves and green stems	occurs in the peripheral regions of stems and leaves.
3. Sclerenchyma (Gk. scleros = hard)	Dead	Sclerenchyma consists of thick walled cells, walls uniformly thick with lignin.	Sclerenchyma is mainly a supporting tissue, which can withstand strains and protect the	• Fibres occur in patches or continuous bands in various parts of stem in
<ul><li>(a) Fibres</li><li>(b) Sclereids</li></ul>	Dead Dead	Walls are thick with lignin. Irregular in shape.	inner thin-walled cells from damage.	<ul> <li>Sclereids occur commonly in fruits and</li> </ul>
		Cell wall very thick making the cell cavity very small.		seeds. Present in some leaves in large numbers.



Fig. 5.2, 5.3, 5.4 Various types of simple tissues

### 5.2.2. Complex tissues

Complex tissues are mainly of two types :

- (i) Xylem
- (ii) Phloem
- Xylem and phloem form a continuous system inside the plants, that is from the roots through the stem and leaves.
- They are known as vascular tissues and form vasular bundles in roots and stems.

**Xylem** (Greek xylo = wood)

- Xylem is a conducting tissue which conducts water and salts upward from roots to leaves.
- Xylem is composed of (a) Tracheids, (b) Vessels (c) Fibres and (d) Xylem Parenchyma (Fig. 5.5)

#### Phloem

- Phloem too is a conducting tissue which conducts the metabolites (food) food synthesised in the leaves to different parts of the plant.
- Phloem is composed of (a) Sieve tube element (b) Companion cells (c) Phloem fibre and (d) Phloem Parenchyma (Fig. 5.6)

The structure, and function of the complex plant tissues is given in Table 5.3.

#### **MODULE - 1** Diversity and Evolution

of Life



 Table 5.3 Structure and function of the components of xylem and phloem

**Tissues and Other Levels of Organization** 

Tissues	Living	Structure	Function
	0r Dood		
	Deau		
Xylem	<b>D</b> 1		
1. Tracheids	Dead	Long cells with pointed ends.	All of them function as a unit to
		Walls thick with lignin.	conduct water and minerals upward
2 37 1		Have pores on the walls	from root to leaves.
2. Vessels	Dead	Cells shorter and broader than	
		trachelds. Walls thick with light	
		and have pores. End walls open	
		and the cens join to form a long	
3 Xylem	Dead	Long cells with very thick lignin	
Fibres	Deau	deposition on the walls no pores	
110105		on the walls	
4. Xylem		Small thin walled cells with	
Parenchyma	Living	cellulose walls.	
Phloem	0		
1. Sieve tube	Living	Elongated sieve elements join to	All of them function as a unit to
		form sieve tubes; cell wall of	translocate food assimilated in the
		cellulose. End walls of the cells	leaves by photosynthesis to different
		have perforations on them, which	parts of the plant.
		give them the name (sieve).	
2. Companion	Living	Long, rectangular cells associated	
cell		with seive cells. Cell wall made	
		of cellulose.	
3. Phloem fibre	Dead	Very long cells with thick lignified	
	<b>.</b>	walls	
4. Phloem	Living	Elongated cells. Cell walls thin	
parenchyma		and made of cellulose.	



**5.2.3 Theories explaining growth of the plant at its shoot apex and root tip** There are two important theories that explain the growth of a plant at the extremities of shoot and root. These are (i) the Tunica corpus theory and (2) the Histogen theory.

#### **Tunica Corpus Theory :**

- Tunica corpus theory was developed for vegetative shoot apex.
- According to this theory, there are two zones of tissues in the apical meristems the tunica (Tunic = cover) consisting of one or more layers of peripheral layers of cells, and the corpus (corpus = body) a mass of cells enclosed by the tunica.
- According to the theory, different planes and rates of cell division and methods of growth in the apex set apart two regions.
- The layers of tunica show anticlinal (perpendicular to periphery) divisions and bring about surface growth.
- In the corpus, cell division is irregular and at various planes resulting in grwoth in volume of the mass.
- Tunica gives rise to the epidermis and cortex. Corpus gives rise to endodermis, pericycle, pith and vascular tissue.

#### **Histogen Theory**

- According to this theory, the apical meristem of stem and root are composed of small mass of cells which are all alike and divide fast (meristematic)
- These meristematic cells form promeristem, which differentiates into three zones dermatogen, periblem and plerome.
- Each every zone consists of a group of initials called a **histogen** (tissue builder).
  - (i) The dermatogen gives rise to epidermis of stems and epiblema of roots.
  - (ii) Periblem (middle layer) gives rise to cortex of stems and roots.
  - (iii) Plerome gives rise to the central meristematic region pericycle, pith and vascular tissue.

#### Classification of plant tissues-at a glance



Diversity and Evolution of Life



Notes 4

## MODULE - 1 Diversity and Evolution

of Life



## INTEXT QUESTIONS 5.2

1. Give Two characteristics and one example of the location of the given tissues in plants in the following table:

1		
- T		_
$\neg$	 	

S

2.

.No.	Tissue	Characteristics	Example of location
(i)	Parenchyma		
(ii)	Collenchyma		
(iii)	Sclerenchyma		
Name	the plant tissues whe	nich	

- (i) conduct water .....
- (ii) conduct food metabolites.

#### **5.3 ANIMAL TISSUES**

As in plants, tissues in animals are also of various types which perform different functions. See the flow-chart given below

Animal	Tissues

Epithetial tissue	<b>Connective Tissue</b>	Muscular tissue	Nervous Tissue
(Protection by covering,	(Binding, support,	(Movement and	(Control and
secretion,	transport)	Locomotion)	co-ordination)
and absoption)			

#### **5.3.1 Epithelial Tissue**

Structural Characteristics : The cells forming epithelial tissue -

- (i) are closely packed with no intercellular spaces in between.
- arise from a non-cellular basement membrane. (ii)
- (iii) are not supplied with blood vessels.

Function : Epithelian tissues line the surfaces, help in absorption, secretion, and also bear protoplasmic projections such as the cilia. (See Table 5.4 and Fig. 5.7)

#### Table 5.4 : Types of epithelial tissue

	Т.''С'	
1. Squamous       Flattened cells with         Epithelium       a centrally placed nucleus.         Have irregular margins.	Lining of air sacs in the lungs. Lining of Kidney tubules. Lining of blood capillaries.	For exchange of $O_2$ and $CO_2$ . For absorption. For exchange of materials.

#### 2. Cubodial Cube like cells with a Lining of salivary For absorption. Epithelium centrally placed nucleus, and pancreatic Cells appear polygonal. ducts. In sweat and For secretion salivary glands. 3. Ciliated Have cilia at Lining of kidney For flow of Notes Epithelium free ends. tubules. nephric filtrate. 4. Columnar Long column-like cells, each Secretion and Lining of stomach, epithelium with nucleus at the basal end instestine absorption 5. Ciliated Cilia at free ends Lining of trachea Flow of fluids in a Columnar particular direction Epithelium 6. Brush bordered Numerous folds at Lining of intestine Increasing the Columnar free ends-folds looking surface area for Epithelium like bristles of a brush. absorption Goblet Cell Simple columnar Central spherical Basement membrane Basement epithelium nucleus Membrane supporting the Cilia goblet cell UUUVARIA Smooth firm outer surface Nucleus

Diversity and Evolution of Life

MODULE - 1



Cytoplasm

Mucus-secreting cell Basement membrane Columnar epitheliurr

Squamous epithelium

Nucleus

Supporting cell

A

Basement membrane

If the epithelial cells are in a single layer, they form simple eptihelium. If the epithelial cells are arranged in many layers, they form compound epithelium or stratified epithelium (many layers). Stratified eptihelium is present in the body, where there is lot of wear and tear. For example the skin and inner lining of cheeks.

#### **BIOLOGY**

0

Fig. 5.7 The structure of different kinds of epithelial tissues

MODULE - 1 **Tissues and Other Levels of Organization** Diversity and Evolution of Life INTEXT QUESTIONS 5.3 1. List the different types of animal tissues Match the items in Column I with those in Column II by writing the 2. Notes corresponding serial number within brackets. Column I Column II **Compound Epithelium** ()(i) Epithelial tissue (a) Basement membrane ()(ii) For increasing the surface area (b)

- (c) Brush-bordered epithelium () (iii) Lining of trachea
- (d) Salivary gland ( ) (iv) Skin
- (e) Ciliated Epithelium () (v) Cuboidal epithelium

### 5.3.2 Connecitve tissue

The connective tissue has two components :

(a) matrix, the ground substance and (b) cells

The matrix and cells are different in different connective tissues (Fig. 5.8).



### A. Proper Connective Tissue

1. Areolar : Most widely spread connective tissue.

The cells forming the tissue are :

- (i) **Fibroblasts**-which form the yellow (elastin) and white (collagen) fibres in the matrix.
- (ii) Macrophages-which help in engulfing bacteria and micro-pathogens.
- (iii) Mast cell-which secretes heparin, that helps in clotting of blood.

**2.** Adipose tissue : It has specialized cells which store fat and provide help in forming paddings.

3. Fibrous : It is mainly made up of fibroblasts. It forms tendons and ligaments.







This is composed of long excitable cells containing parallel microfilaments of contractile proteins, as in actin, myosin, troponin and tropomysin. Because of its elongated shape, muscle cell is called a muscle fibre. The muscle fibres of vertebrates are of three different types (i) Striated (ii) Unstriated and (iii) Cardiac (Fig. 5.9) according to the shape and functions as mentioned in Table 5.5 and Fig. 5.9.

Striated/Voluntary/Skeletal	Unstriated/	Cardiac
	Involuntary	
Location		
1. Attached to the skeleton like	In the walls of body organs	Walls of heart.
head, limbs, face etc.	like stomach, intestines.	
Shape		
Elongated, cylindrical, unbranched fibres	Spindle shaped, tapering.	Elongated, cylindrical, branched.
Myofibrils so arranged in the	No such striations seen as	Striations (stripes) seen.
cytoplasm, that there are	myofibrils are not uniformly	_
striations seen.	arranged .	
Sarcolemma		
Thin and tough membrane	Thin cell membrane, no	Thin
sarcolemma of the fibre (cell).	sarcolemma.	
Nucleus		
Multi nucleated, Peripheral	Uninucleated, centrally	One nucleus in each unit,
nuclei.	placed.	centrally placed.
Blood Supply		
Rich	Poor	Rich
Intercalate Discs		
Absent	Absent	Present
Voluntary (Contracts at will)	Involuntary	Involuntary

#### Table 5.5 Types of Muscle Fibres

MODULE - 1 Diversity and Evolution of Life







Fig. 5.9 Types of Vertebrate Muscle Tissue

## MODULE - 1

Diversity and Evolution of Life

The muscle fibres have the following characteristics:

- (i) Excitability, (respond to stimulus)
- Extensibility, (stretch) (ii)
- Contractility, (contract) (iii)
- (iv) Elasticity, (move back to the original position)

## INTEXT QUESTIONS 5.4

- Name the different types of cells found in the different types of connective tissue. 1.
- ..... 2. Match the item in Column I with those in Column II, by writing the corresponding serial number within brackets:

0110	sponding serial name	<b>U</b> I 111		
	Column I			Column II
a.	Unstripped muscles	()	(i)	multinucleate
b.	Myofibrils	()	(ii)	run parallel to each other in a striped
				muscle
c.	Sarcolemma	()	(iii)	cardiac muscles
d.	Striped muscle	()	(iv)	outer tough membrane of a striped muscle
				fibre
e.	Branched myofibrils	( )	(v)	involuntary
	lervous Tissues			

#### 5.3.4

Nervous tissues has two kinds of cells i.e. neurons and neuroglia cells

#### Neurons

Neuron is the functional unit of nervous tissue. Neurons are also called nerve cells. Nervous tissues constitute the brain, spinal cord, nerves and the sensory cells and sense organs.

A single neuron has a generalised appearance as shown in the Fig. 5.10.





Like any other cell of the body, the nerve cell or neuron has the main cell body called **cyton** from which project out a varying number of processes –one of which is usually very long. This long fibre is called the **axon**.

The smaller branching processes of the cyton are called the **dendrites** (GK dendros = tree). The cells bounded by plasma membrane, possess a nucleus and other organelles like mitochondria.

The cyton also contains dark granules called **Nissi** bodies. These are made of RNA and Protein.

**Transmission of nerve impulse** – The branching dendrites receive the stimulus and transmit it through the cyton to the axon, which finally transmits it through its variously branched ends into either a muscle (to order it to contract) or to a gland (to order it to secrete). The axon constitutes the nerve fibre. The nerve fibre may or may not be covered by an extra sheath called **medullary sheath** secreted by sheath cells. It is made of **myelin** a lipid like substance. Accordingly, the nerve fibre is termed **medullated** and **non-medullated**. The medullary sheath is not continuous and is broken at **nodes of Ranvier** (Fig. 5.10).



1. What is the function of the nervous tissue?

.....

2. What is the direction of the "flow of impulse" within a nerve cell from its dendrites to its axon end or from its axon end toward its dendrites?

.....

- 3. What are the following parts in a nerve cell?
  - (i) Cyton.....
  - (ii) Dendrite.....
  - (iii) Axon.....
  - (iv) Medullary sheath.....
  - (v) Node of Ranvier .....

#### 5.4 LEVELS OF ORGANISATION – CELL TO ORGANISM

We started the lesson by talking about the smallest unit of life in any living organism i.e. the cell. The cell has a very complex system of its organelles, each organelle concerned with a particular task or activity, and each activity contributing to the total performance of the cell. Thus there is a division of labour at the cellular level. As evolution progressed and larger and larger organisms appeared with enormous number of cells in the body, it became necessary that the bodily functions are distributed among different groups of cells or tissues even among groups of tissues. Such higher and higher stages or groupings are known as the levels of organization. These levels are as follows:

**BIOLOGY** 



MODULE - 1



### **MODULE - 1** Diversity and Evolution

of Life



Notes

- (i) **Cellular Levels of Organization** The organization of the activities by different organelles in a single cell. Example, white blood cells or a green cells of a leaf.
- (ii) **Tissue Level** The aggregates of cells of same origin and having same function, example, the surface epithelium of our skin or the dividing cells at the root cap of a plant.
- (iii) **Tissue System** Generally seen in plants where two or more different cell types combine to perform a particular activity. Example Vascular tissue e.g. veins of a leaf, consisting of xylem and phloem, for transport of water and food materials.
- (iv) **Organ Level–** A distinct recognizable part of the body, composed of a variety of tissues and performing one or more special functions which contribute to the well being of the organism. Example : Liver in animals and leaf in plants.
- (v) **Organ System-** Combination of a set of organs all of which are usually devoted to one general function. Example : respiratory system (consisting of lungs, trachea, and diaphragm) in man or the shoot system (consisting of leaves, stem and branches) in a plant.
- (vi) **Organism–** The complete individual made of different organ systems. Examples: man, monkey, or a mustard plant.

## INTEXT QUESTIONS 5.6

1. Rearrange the following levels of organizations in their correct sequences:tissue, cell, organ, organism, organ system.

.....

2. Complete the following Table by giving one example of each of the following in an animal and plant.

Level of	Examp	Examples		
Organisation	Animal	Plant		
Cell				
Tissue				
Organ				
Organ system				
Organism				



• A tissue is a group of cells which are essentially of the same kind and of the same origin and performing similar function.

- In plants there are, first of all two major categories of tissues- meristematic (dividing and undifferentiated) and permanent (specialized) tissues.
- Meristematic tissue is located at all growth points.
- Permanent tissue consists of the simple tissue (parenchyma, collenchyma and sclerenchyma) and complex tissue (xylem and phloem).
- The animal tissues consist of epithelium (closely packed cells usually on surfaces,) connective tissue which primarily support, connect or bind the body parts together (bones blood etc.), the contractile muscular tissue (different muscles,) and nervous tissue consisting of nerve cells adapted for conducting the message (brain cells,)
- The various tissues in both plants and animals are grouped together to form an organ. The different organs together form the organ system and the various organs systems together constitute the organism or the individual. Thus there are different levels of organization with increasing complexity and specialization from cell to organism.

# TERMINAL EXERCISES

- 1. What is a tissue?
- 2. State one main structural characteristic and the special activity of the following tissue:

meristem, sclerenchyma, xylem, phloem, epithelium, muscle, nervous tissue.

- 3. In what way do the following tissues differ from the one stated:-
  - (i) Connective tissue from epithelial tissue
  - (ii) Bone from blood
  - (iii) Phloem from xylem
  - (iv) Squamous epithelium from columnar epithelium
  - (v) Tracheids from wood fibres
- 4. Name the different levels of organizations in animals (such as humans) giving one example of each.



## ANSWERS TO INTEXT QUESTIONS

- 5.1 1. a group of cells with similar origin, structure and function
  - 2. (i) Merisematic;
    - (ii) Lateral meristem
    - (iii) Simple
    - (iv) Permanent
  - 3. arising from same embryonic layer of cells

#### BIOLOGY

Diversity and Evolution of Life

MODULE - 1



Diversity and Evolution of Life 4. histology	
4. Instology	
5. composed of more than one type of cells all cooperating in	performing
common function	r8
<b>5.2</b> 1. S.No. Tissue Characteristics Example of locat	ion
1. Parenchyma 1. Round cells 1. Root, stem	and leaves
Notes2. Living2. Collenchyma1. Polygonal cells1. Petiole and	Mid-rib of
with thickening leaves	
2. Living	
3. Sclerenchyma 1. Elongated or 1. Woody Sten	18
irregular	
in shape 2 Dead and thick	
walled	
2. xylem, phloem	
<b>5.3</b> 1. Epithelial, connective, muscular, nervous	
2. a-iv, b-i, c-ii, e-iii	
<b>5.4</b> 1. Firbroblasts - areolar	
Macrophages - areolar	
Mast cells - areolar	
Cartilage cells/chondrocyte - chondrocyte-cartilage	
Bone cells/osteocyte - bone	
Blood cells/WBC RBC - blood	
2. a (v); (b) (ii); c (iv); d (i); e. (iii)	
<b>5.5</b> 1. sensory	
2. Dendrite to the axon	
<ul><li>3. (i) cell-body (ii) thin processes of cyton (iii) set (iv) medullary layer (v) interruptions in medullary sheath</li></ul>	nsory fibre
<b>5.6</b> 1. Cell, tissue, organ, organ system, organism	
2. refer to text subsection 5.4	

## **MODULE - II**

## FORMS AND FUNCTIONS OF PLANTS AND ANIMALS

- 06 Root system
- 07 Shoot system
- 08 Absorption, Transport and Water Loss in Plants
- 09 Nutrition in plants Mineral Nutrition
- 10 Nitrogen Metabolism
- 11 Photosynthesis
- 12 Respiration in Plants
- 13 Nutrition and Digestion
- 14 Respiration and Elimination of Nitrogenous Wastes
- 15 Circulation of Body Fluids
- 16 Locomotion and Movement
- 17 Coordination and Control The Nervous and Endocrine Systems
- 18 Homeostasis: The Steady State