



## ARITHMETIC AND GEOMETRIC PROGRESSIONS

Succession of numbers of which one number is designated as the first, other as the second, another as the third and so on gives rise to what is called a sequence. Sequences have wide applications. In this lesson we shall discuss particular types of sequences called arithmetic sequence, geometric sequence and also find arithmetic mean (A.M), geometric mean (G.M) between two given numbers. We will also establish the relation between A.M and G.M.

**Let us consider the following problems :**

- (a) A man places a pair of newly born rabbits into a warren and wants to know how many rabbits he would have over a certain period of time. A pair of rabbits will start producing offsprings two months after they were born and every following month one new pair of rabbits will appear. At the beginning the man will have in his warren only one pair of rabbits, during the second month he will have the same pair of rabbits, during the third month the number of pairs of rabbits in the warren will grow to two; during the fourth month there will be three pairs of rabbits in the warren. Thus, the number of pairs of rabbits in the consecutive months are :

1, 1, 2, 3, 5, 8, 13, ...

- (b) The recurring decimal  $0.\bar{3}$  can be written as a sum

$$0.\bar{3} = 0.3 + 0.03 + 0.003 + 0.0003 \dots$$

- (c) A man earns Rs.10 on the first day, Rs. 30 on the second day, Rs. 50 on the third day and so on. The day to day earning of the man may be written as 10, 30, 50, 70, 90, ...

We may ask what his earnings will be on the 10<sup>th</sup> day in a specific month.

Again let us consider the following sequences:

MODULE - III  
Sequences And  
Series



Notes

$$(1) \quad 2, 4, 8, 16, \dots \quad (2) \quad \frac{1}{9}, -\frac{1}{27}, \frac{1}{81}, -\frac{1}{243}, \dots$$

$$(3) \quad 0.01, 0.0001, 0.000001, \dots$$

In these three sequences, each term except the first, progresses in a definite order but different from the order of other three problems. In this lesson we discuss those sequences whose term progresses in a definite order.



### OBJECTIVES

After studying this lesson, you will be able to :

- describe the concept of a sequence (progression);
- define an A.P. and cite examples;
- find common difference and general term of a A.P;
- find the fourth quantity of an A.P. given any three of the quantities  $a$ ,  $d$ ,  $n$  and  $t_n$ ;
- calculate the common difference or any other term of the A.P. given any two terms of the A.P;
- derive the formula for the sum of 'n' terms of an A.P;
- calculate the fourth quantity of an A.P. given three of  $S$ ,  $n$ ,  $a$  and  $d$ ;
- insert A.M. between two numbers;
- solve problems of daily life using concept of an A.P;
- state that a geometric progression is a sequence increasing or decreasing by a definite multiple of a non-zero number other than one;
- identify G.P.'s from a given set of progressions;
- find the common ratio and general term of a G.P;
- calculate the fourth quantity of a G.P when any three of the quantities  $t_n$ ,  $a$ ,  $r$  and  $n$  are given;
- calculate the common ratio and any term when two of the terms of the G.P. are given;
- write progression when the general term is given;
- derive the formula for sum of  $n$  terms of a G.P;
- calculate the fourth quantity of a G.P. if any three of  $a$ ,  $r$ ,  $n$  and  $S$  are given;
- derive the formula for sum ( $S_\infty$ ) of infinite number of terms of a G.P. when  $|r| < 1$ ;
- find the third quantity when any two of  $S_\infty$ ,  $a$  and  $r$  are given;



- convert recurring decimals to fractions using G.P;
- insert G.M. between two numbers; and
- establish relationship between A.M. and G.M.

### EXPECTED BACKGROUND KNOWLEDGE

- Laws of indices
- Simultaneous equations with two unknowns.
- Quadratic Equations.

### 13.1 SEQUENCE

A sequence is a collection of numbers specified in a definite order by some assigned law, whereby a definite number  $a_n$  of the set can be associated with the corresponding positive integer  $n$ .

The different notations used for a sequence are.

1.  $a_1, a_2, a_3, \dots, a_n, \dots$
2.  $a_n, n = 1, 2, 3, \dots$
3.  $\{a_n\}$

Let us consider the following sequences :

1.  $1, 2, 4, 8, 16, 32, \dots$
2.  $1, 4, 9, 16, 25, \dots$
3.  $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \dots$
4.  $1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \frac{1}{6}, \dots$

In the above examples, the expression for  $n^{\text{th}}$  term of the sequences are as given below :

$$(1) a_n = 2^{n-1} \quad (2) a_n = n^2 \quad (3) a_n = \frac{n}{n+1} \quad (4) a_n = \frac{1}{n}$$

for all positive integer  $n$ .

Also for the first problem in the introduction, the terms can be obtained from the relation

$$a_1 = 1, a_2 = 1, a_n = a_{n-2} + a_{n-1}, n \geq 3$$

A finite sequence has a finite number of terms. An infinite sequence contains an infinite number of terms.

### 13.2 ARITHMETIC PROGRESSION

Let us consider the following examples of sequence, of numbers :

- (1)  $2, 4, 6, 8, \dots$
- (2)  $1, \frac{3}{2}, 2, \frac{5}{2}, \dots$

MODULE - III  
Sequences And  
Series



## Notes

$$(3) \quad 10, 8, 6, 4, \dots \quad (4) \quad -\frac{1}{2}, -1, -\frac{3}{2}, -2, -\frac{5}{2}, \dots$$

Note that in the above four sequences of numbers, the first terms are respectively 2, 1, 10, and  $-\frac{1}{2}$ . The first term has an important role in this lesson. Also every following term of the sequence has certain relation with the first term. What is the relation of the terms with the first term in Example (1) ?

$$\text{First term} = 2$$

$$\text{Second term} = 4 = 2 + 1 \times 2$$

$$\text{Third term} = 6 = 2 + 2 \times 2$$

$$\text{Fourth term} = 8 = 2 + 3 \times 2$$

and so on.

The consecutive terms in the above sequence are obtained by adding 2 to its preceding term. i.e., the difference between any two consecutive terms is the same.

A finite sequence of numbers with this property is called an arithmetic progression.

A sequence of numbers with finite terms in which the difference between two consecutive terms is the same non-zero number is called the Arithmetic Progression or simply A. P.

The difference between two consecutive terms is called the common difference of the A. P. and is denoted by ' $d$ '.

In general, an A. P. whose first term is  $a$  and common difference is  $d$  is written as  $a, a + d, a + 2d, a + 3d, \dots$

Also we use  $t_n$  to denote the  $n$ th term of the progression.

### 13.2.1 GENERAL TERM OF AN A. P.

Let us consider A. P.

$$a, a + d, a + 2d, a + 3d, \dots$$

Here, first term  $(t_1) = a$

$$\text{second term } (t_2) = a + d = a + (2 - 1) d,$$

$$\text{third term } (t_3) = a + 2d = a + (3 - 1) d$$

By observing the above pattern,  $n$ th term can be written as:

$$t_n = a + (n - 1) d$$

Hence, if the first term and the common difference of an A. P. are known then any term of A. P. can be determined by the above formula.

Sometimes, the  $n$ th term of an A. P. is expressed, in terms of  $n$ , e.g.  $t_n = 2n - 1$ . In that case, the



A. P. will be obtained by substituting  $n = 1, 2, 3, \dots$  in the expression. In this case, the terms of the A. P. are 1, 3, 5, 7, 9, ...

**Note.:**

- (i) If the same non-zero number is added to each term of an A. P. the resulting sequence is again an A. P.
- (ii) If each term of an A. P. is multiplied by the same non-zero number, the resulting sequence is again an A. P.

**Example 13.1** Find the 10<sup>th</sup> term of the A. P.: 2, 4, 6, ...

**Solution :** Here the first term ( $a$ ) = 2 and common difference  $d = 4 - 2 = 2$

Using the formula  $t_n = a + (n - 1) d$ , we have

$$t_{10} = 2 + (10 - 1) 2 = 2 + 18 = 20$$

Hence, the 10th term of the given A. P. is 20.

**Example 13.2** The 10<sup>th</sup> term of an A. P. is  $-15$  and 31<sup>st</sup> term is  $-57$ , find the 15<sup>th</sup> term.

**Solution :** Let  $a$  be the first term and  $d$  be the common difference of the A. P. Then from the formula:

$$t_n = a + (n - 1) d, \text{ we have}$$

$$t_{10} = a + (10 - 1) d = a + 9d$$

$$t_{31} = a + (31 - 1) d = a + 30d$$

We have,

$$a + 9d = -15 \quad \dots(1)$$

$$a + 30d = -57 \quad \dots(2)$$

Solve equations (1) and (2) to get the values of  $a$  and  $d$ .

Subtracting (1) from (2), we have

$$21d = -57 + 15 = -42$$

$$\therefore d = \frac{-42}{21} = -2$$

Again from (1),  $a = -15 - 9d = -15 - 9(-2) = -15 + 18 = 3$

Now  $t_{15} = a + (15 - 1)d$

$$= 3 + 14(-2) = -25$$

**Example 13.3** Which term of the A. P.: 5, 11, 17, ... is 119 ?

**Solution :** Here  $a = 5$ ,  $d = 11 - 5 = 6$

MODULE - III  
Sequences And  
Series

Notes

$$t_n = 119$$

We know that

$$t_n = a + (n - 1) d$$

$$\Rightarrow 119 = 5 + (n - 1) \times 6$$

$$\Rightarrow (n - 1) = \frac{119 - 5}{6} = 19$$

$$\therefore n = 20$$

Therefore, 119 is the 20th term of the given A. P.

**Example 13.4** Is 600 a term of the A. P.: 2, 9, 16, ...?

**Solution :** Here,  $a = 2$ , and  $d = 9 - 2 = 7$ .

Let 600 be the  $n^{\text{th}}$  term of the A. P. We have  $t_n = 2 + (n - 1) 7$

According to the question,

$$2 + (n - 1) 7 = 600$$

$$\therefore (n - 1) 7 = 598$$

$$\text{or } n = \frac{598}{7} + 1$$

$$\therefore n = 86\frac{3}{7}$$

Since  $n$  is a fraction, it cannot be a term of the given A. P. Hence, 600 is not a term of the given A. P.

**Example 13.5** The common difference of an A. P. is 3 and the 15<sup>th</sup> term is 37. Find the first term.

**Solution :** Here,  $d = 3$ ,  $t_{15} = 37$ , and  $n = 15$

Let the first term be  $a$ . We have

$$t_n = a + (n - 1) d$$

$$37 = a + (15 - 1) 3$$

$$\text{or, } 37 = a + 42$$

$$\therefore a = -5$$

Thus, first term of the given A. P. is  $-5$ .



**Example 13.6** If  $a + b + c = 0$  and  $\frac{a}{b+c}, \frac{b}{c+a}, \frac{c}{a+b}$  are in A. P., then prove that

$\frac{1}{b+c}, \frac{1}{c+a}, \frac{1}{a+b}$  are also in A. P.

**Solution.** : Since  $\frac{a}{b+c}, \frac{b}{c+a}, \frac{c}{a+b}$  are in A. P., therefore

$$\frac{b}{c+a} - \frac{a}{b+c} = \frac{c}{a+b} - \frac{b}{c+a}$$

or,  $\left(\frac{b}{c+a} + 1\right) - \left(\frac{a}{b+c} + 1\right) = \left(\frac{c}{a+b} + 1\right) - \left(\frac{b}{c+a} + 1\right)$

or,  $\frac{a+b+c}{c+a} - \frac{a+b+c}{b+c} = \frac{a+b+c}{a+b} - \frac{a+b+c}{c+a}$

or,  $\frac{1}{c+a} - \frac{1}{b+c} = \frac{1}{a+b} - \frac{1}{c+a}$  (Since  $a + b + c \neq 0$ )

or,  $\frac{1}{b+c}, \frac{1}{c+a}, \frac{1}{a+b}$  are in A. P.



### CHECK YOUR PROGRESS 13.1

- Find the  $n^{\text{th}}$  term of each of the following A. P's. :  
(a) 1, 3, 5, 7, ...      (b) 3, 5, 7, 9, ...
- If  $t_n = 2n + 1$ , then find the A. P.
- Which term of the A. P.  $2\frac{1}{2}, 4, 5\frac{1}{2}, \dots$  is 31? Find also the 10<sup>th</sup> term?
- Is  $-292$  a term of the A. P. 7, 4, 1,  $-2, \dots$ ?
- The  $m^{\text{th}}$  term of an A. P. is  $n$  and the  $n^{\text{th}}$  term is  $m$ . Show that its  $(m + n)^{\text{th}}$  term is zero.
- Three numbers are in A. P. The difference between the first and the last is 8 and the product of these two is 20. Find the numbers.
- The  $n^{\text{th}}$  term of a sequence is  $na + b$ . Prove that the sequence is an A. P. with common difference  $a$ .



Notes

### 13.3 TO FIND THE SUM OF FIRST $n$ TERMS IN AN A. P.

Let  $a$  be the first term and  $d$  be the common difference of an A. P. Let  $l$  denote the last term, i.e., the  $n^{\text{th}}$  term of the A. P. Then,

$$l = t_n = a + (n - 1)d \quad \dots \text{(i)}$$

Let  $S_n$  denote the sum of the first  $n$  terms of the A. P. Then

$$S_n = a + (a + d) + (a + 2d) + \dots + (l - 2d) + (l - d) + l \quad \dots \text{(ii)}$$

Reversing the order of terms in the R. H. S. of the above equation, we have

$$S_n = l + (l - d) + (l - 2d) + \dots + (a + 2d) + (a + d) + a \quad \dots \text{(iii)}$$

Adding (ii) and (iii) vertically, we get

$$2S_n = (a + l) + (a + l) + (a + l) + \dots \text{containing } n \text{ terms} = n(a + l)$$

$$\text{i.e., } S_n = \frac{n}{2}(a + l)$$

$$\text{Also } S_n = \frac{n}{2}[2a + (n - 1)d] \quad [\text{From (i)}]$$

It is obvious that  $t_n = S_n - S_{n-1}$

**Example 13.7** Find the sum of  $2 + 4 + 6 + \dots n$  terms.

**Solution.:** Here  $a = 2$ ,  $d = 4 - 2 = 2$

Using the formula  $S_n = \frac{n}{2}[2a + (n - 1)d]$ , we get

$$\begin{aligned} S_n &= \frac{n}{2}[2 \times 2 + (n - 1)2] \\ &= \frac{n}{2}[2 + 2n] = \frac{2n(n + 1)}{2} \\ &= n(n + 1) \end{aligned}$$

**Example 13.8** Find the sum of the sequence  $2, 3, 5, 9, 8, 15, 11, \dots$  to  $(2n + 1)$  terms

**Solution.** Let  $S$  denote the sum. Then

$$\begin{aligned} S &= 2 + 3 + 5 + 9 + 8 + 15 + 11 + \dots (2n + 1) \text{ terms} \\ &= [2 + 5 + 8 + 11 + \dots (n + 1) \text{ terms}] + [3 + 9 + 15 + 21 + \dots n \text{ terms}] \\ &= \frac{n + 1}{2}[2 \times 2 + (n + 1 - 1)3] + \frac{n}{2}[2 \times 3 + (n - 1)6] \end{aligned}$$





$$\begin{aligned}
 &= \frac{n+1}{2} [4 + 3n] + \frac{n}{2} [6 + 6n - 6] \\
 &= \frac{(n+1)(3n+4)}{2} + 3n^2 \\
 &= \frac{1}{2} [3n^2 + 7n + 4 + 6n^2] \\
 &= \frac{1}{2} [9n^2 + 7n + 4]
 \end{aligned}$$

**Example 13.9** The 35<sup>th</sup> term of an A. P. is 69. Find the sum of its 69 terms.

**Solution.** Let  $a$  be the first term and  $d$  be the common difference of the A. P.

We have

$$t_{35} = a + (35 - 1)d = a + 34d.$$

$$\therefore a + 34d = 69 \quad \dots (i)$$

Now by the formula,  $S_n = \frac{n}{2} [2a + (n - 1)d]$

We have

$$\begin{aligned}
 S_{69} &= \frac{69}{2} [2a + (69 - 1)d] \\
 &= 69(a + 34d) \quad \text{[using (i)]} \\
 &= 69 \times 69 = 4761
 \end{aligned}$$

**Example 13.10** The first term of an A. P. is 10, the last term is 50. If the sum of all the terms is 480, find the common difference and the number of terms.

**Solution :** We have:  $a = 10, l = t_n = 50, S_n = 480.$

By substituting the values of  $a, t_n$  and  $S_n$  in the formulae

$$S_n = \frac{n}{2} [2a + (n - 1)d] \text{ and } t_n = a + (n - 1)d, \text{ we get}$$

$$480 = \frac{n}{2} [20 + (n - 1)d] \quad \dots (i)$$

$$50 = 10 + (n - 1)d \quad \dots (ii)$$

$$\text{From (ii), } (n - 1)d = 50 - 10 = 40 \quad \dots (iii)$$

MODULE - III  
Sequences And  
Series

Notes

From (i), we have

$$480 = \frac{n}{2} (20 + 40) \quad \text{using (i)}$$

$$\text{or, } 60n = 2 \times 480$$

$$\therefore n = \frac{2 \times 480}{60} = 16$$

From (iii),

$$\therefore d = \frac{40}{15} = \frac{8}{3} \quad (\text{as } n - 1 = 16 - 1 = 15)$$

**Example 13.11** Let the  $n^{\text{th}}$  term and the sum of  $n$  terms of an A. P. be  $p$  and  $q$  respectively.

Prove that its first term is  $\left(\frac{2q - pn}{n}\right)$ .

**Solution:** In this case,  $t_n = p$  and  $S_n = q$

Let  $a$  be the first term of the A. P.

$$\text{Now, } S_n = \frac{n}{2}(a + t_n)$$

$$\text{or, } \frac{n}{2}(a + p) = q$$

$$\text{or, } a + p = \frac{2q}{n}$$

$$\text{or, } a = \frac{2q}{n} - p$$

$$\therefore a = \frac{2q - pn}{n}$$

**CHECK YOUR PROGRESS 13.2**

- Find the sum of the following A. P.'s.
  - 8, 11, 14, 17, ... up to 15 terms
  - 8, 3, -2, -7, -12, ... up to  $n$  terms.
- How many terms of the A. P.: 27, 23, 19, 15, ... have a sum 95?
- A man takes an interest-free loan of Rs. 1740 from his friend agreeing to repay in monthly



instalments. He gives Rs. 200 in the first month and diminishes his monthly instalments by Rs. 10 each month. How many months will it take to repay the loan?

4. How many terms of the progression 3, 6, 9, 12, ... must be taken at the least to have a sum not less than 2000?
5. In a children potato race,  $n$  potatoes are placed 1 metre apart in a straight line. A competitor starts from a point in the line which is 5 metre from the nearest potato. Find an expression for the total distance run in collecting the potatoes, one at a time and bringing them back one at a time to the starting point. Calculate the value  $n$  if the total distance run is 162 metres.
6. If the sum of first  $n$  terms of  $a$  sequence be  $an^2 + bn$ , prove that the sequence is an A. P. and find its common difference?

### 13.4 ARITHMETIC MEAN (A. M.)

When three numbers  $a, A$  and  $b$  are in A. P., then  $A$  is called the arithmetic mean of numbers  $a$  and  $b$ . We have,  $A - a = b - A$

$$\text{or, } A = \frac{a+b}{2}$$

Thus, the required A. M. of two numbers  $a$  and  $b$  is  $\frac{a+b}{2}$ . Consider the following A. P. :

3, 8, 13, 18, 23, 28, 33.

There are five terms between the first term 3 and the last term 33. These terms are called *arithmetic means* between 3 and 33. Consider another A. P. : 3, 13, 23, 33. In this case there are two arithmetic means 13, and 23 between 3 and 33.

Generally any number of arithmetic means can be inserted between any two numbers  $a$  and  $b$ . Let  $A_1, A_2, A_3, \dots, A_n$  be  $n$  arithmetic means between  $a$  and  $b$ , then.

$a, A_1, A_2, A_3, \dots, A_n, b$  is an A. P.

Let  $d$  be the common difference of this A. P. Clearly it contains  $(n + 2)$  terms

$$\begin{aligned} \therefore b &= (n + 2)^{\text{th}} \text{ term} \\ &= a + (n + 1) d \end{aligned}$$

$$\therefore d = \frac{b-a}{n+1}$$

$$\text{Now, } A_1 = a + d \Rightarrow A_1 = \left( a + \frac{b-a}{n+1} \right) \quad \dots(i)$$

MODULE - III  
Sequences And  
Series

Notes

$$A_2 = a + 2d \Rightarrow A_2 = \left( a + \frac{2(b-a)}{n+1} \right) \quad \dots(ii)$$

$$\vdots$$

$$A_n = a + nd \Rightarrow A_n = \left( a + \frac{n(b-a)}{n+1} \right) \quad \dots (n)$$

These are required  $n$  arithmetic means between  $a$  and  $b$ .

Adding (i), (ii), ..., (n), we get

$$\begin{aligned} A_1 + A_2 + \dots + A_n &= na + \dots + \frac{b-a}{n+1} [1+2+\dots+n] \\ &= na + \left( \frac{b-a}{n+1} \right) \left( \frac{n(n+1)}{2} \right) = na + \frac{n(b-a)}{2} = \frac{n(a+b)}{2} \\ &= n [\text{Single A. M. between } a \text{ and } b] \end{aligned}$$

**Example 13.12** Insert five arithmetic means between 8 and 26.

**Solution :** Let  $A_1, A_2, A_3, A_4$  and  $A_5$  be five arithmetic means between 8 and 26.

Therefore, 8,  $A_1, A_2, A_3, A_4, A_5, 26$  are in A. P. with  $a = 8, b = 26, n = 7$

We have  $26 = 8 + (7 - 1) d$

$$\therefore d = 3$$

$$\therefore A_1 = a + d = 8 + 3 = 11, A_2 = a + 2d = 8 + 2 \times 3 = 14$$

$$A_3 = a + 3d = 17, A_4 = a + 4d = 20$$

$$A_5 = a + 5d = 23$$

Hence, the five arithmetic means between 8 and 26 are 11, 14, 17, 20 and 23.

**Example 13.13** The ' $n$ ', A. M's between 20 and 80 are such that the ratio of the first mean and the last mean is 1 : 3. Find the value of  $n$ .

**Solution :** Here, 80 is the  $(n+2)^{\text{th}}$  term of the A. P., whose first term is 20. Let  $d$  be the common difference.

$$\therefore 80 = 20 + (n+2-1) d$$

$$\text{or, } 80 - 20 = (n+1) d$$

$$\text{or, } d = \frac{60}{n+1}$$

$$\text{The first A. M.} = 20 + \frac{60}{n+1} = \frac{20n + 20 + 60}{n+1}$$



$$= \frac{20n + 80}{n + 1}$$

$$\text{The last A. M.} = 20 + n \times \frac{60}{n + 1} = \frac{80n + 20}{n + 1}$$

We have

$$\frac{20n + 80}{n + 1} : \frac{80n + 20}{n + 1} = 1 : 3$$

$$\text{or, } \frac{n + 4}{4n + 1} = \frac{1}{3}$$

$$\text{or, } 4n + 1 = 3n + 12$$

$$\text{or, } n = 11$$

∴ The number of A. M's between 20 and 80 is 11.

**Example 13.14** If  $x, y, z$  are in A. P., show that

$$(x + 2y - z)(2y + z - x)(z + x - y) = 4xyz$$

**Solution :**  $x, y, z$  are in A. P.

$$\therefore y = \frac{x + z}{2} \quad \dots(i)$$

$$\begin{aligned} \therefore (x + 2y - z)(2y + z - x)(z + x - y) & \\ = (x + x + z - z)(x + z + z - x)(2y - y) & \quad \text{[From (i)]} \\ = (2x)(2z)(y) = 4xyz & \\ = \text{R. H. S.} & \end{aligned}$$



### CHECK YOUR PROGRESS 13.3

1. Prove that if the number of terms of an A. P. is odd then the middle term is the A. M. between the first and last terms.
2. Between 7 and 85,  $m$  number of arithmetic means are inserted so that the ratio of  $(m - 3)^{\text{th}}$  and  $m^{\text{th}}$  means is 11 : 24. Find the value of  $m$ .
3. Prove that the sum of  $n$  arithmetic means between two numbers is  $n$  times the single A. M. between them.
4. If the A. M. between  $p^{\text{th}}$  and  $q^{\text{th}}$  terms of an A. P., be equal and to the A. M. between  $r^{\text{th}}$  and  $s^{\text{th}}$  terms of the A. P., then show that  $p + q = r + s$ .



Notes

## 13.5 GEOMETRICAL PROGRESSION

Let us consider the following sequence of numbers :

$$(1) \quad 1, 2, 4, 8, 16, \dots \quad (2) \quad 3, 1, \frac{1}{3}, \frac{1}{9}, \frac{1}{27}, \dots$$

$$(3) \quad 1, -3, 9, -27, \dots \quad (4) \quad x, x^2, x^3, x^4, \dots$$

If we see the patterns of the terms of every sequence in the above examples each term is related to the leading term by a definite rule.

For Example (1), the first term is 1, the second term is twice the first term, the third term is  $2^2$  times of the leading term.

Again for Example (2), the first term is 3, the second term is  $\frac{1}{3}$  times of the first term, third term is  $\frac{1}{3^2}$  times of the first term.

A sequence with this property is called a *geometric progression*.

A sequence of numbers in which the ratio of any term to the term which immediately precedes is the same non zero number (other than 1), is called a geometric progression or simply G. P. This ratio is called the common ratio.

Thus,  $\frac{\text{Second term}}{\text{First term}} = \frac{\text{Third term}}{\text{Second term}} = \dots$  is called the common ratio of the geometric progression.

Examples (1) to (4) are geometric progressions with the first term 1, 3, 1,  $x$  and with common ratio  $2, \frac{1}{3}, -3$ , and  $x$  respectively.

The most general form of a G. P. with the first term  $a$  and common ratio  $r$  is  $a, ar, ar^2, ar^3, \dots$

### 13.5.1 GENERAL TERM

Let us consider a geometric progression with the first term  $a$  and common ratio  $r$ . Then its terms are given by

$$a, ar, ar^2, ar^3, \dots$$

In this case,  $t_1 = a = ar^{1-1}$

$$t_2 = ar = ar^{2-1}$$

$$t_3 = ar^2 = ar^{3-1}$$

$$t_4 = ar^3 = ar^{4-1}$$



Notes

... ..

On generalisation, we get the expression for the  $n^{\text{th}}$  term as

$$t_n = ar^{n-1} \quad \dots \text{(A)}$$

### 13.5.2 SOME PROPERTIES OF A G. P.

- (i) If all the terms of a G. P. are multiplied by the same non-zero quantity, the resulting series is also in G. P. The resulting G. P. has the same common ratio as the original one.

If  $a, b, c, d, \dots$  are in G. P.

then  $ak, bk, ck, dk \dots$  are also in G. P.  $(k \neq 0)$

- (ii) If all the terms of a G. P. are raised to the same power, the resulting series is also in G. P.

Let  $a, b, c, d \dots$  are in G. P.

the  $a^k, b^k, c^k, d^k, \dots$  are also in G. P.  $(k \neq 0)$

The common ratio of the resulting G. P. will be obtained by raising the same power to the original common ratio.

**Example 13.15** Find the 6<sup>th</sup> term of the G. P.: 4, 8, 16, ...

**Solution :** In this case the first term  $(a) = 4$

Common ratio  $(r) = 8 \div 4 = 2$

Now using the formula  $t_n = ar^{n-1}$ , we get

$$t_6 = 4 \times 2^{6-1} = 4 \times 32 = 128$$

Hence, the 6<sup>th</sup> term of the G. P. is 128.

**Example 13.16** The 4<sup>th</sup> and the 9<sup>th</sup> term of a G. P. are 8 and 256 respectively. Find the G. P.

**Solution :** Let  $a$  be the first term and  $r$  be the common ratio of the G. P., then

$$t_4 = ar^{4-1} = ar^3$$

$$t_9 = ar^{9-1} = ar^8$$

According to the question,  $ar^8 = 256 \quad \dots \text{(1)}$

and  $ar^3 = 8 \quad \dots \text{(2)}$

$$\therefore \frac{ar^8}{ar^3} = \frac{256}{8}$$

or  $r^5 = 32 = 2^5$

$$\therefore r = 2$$

Again from (2),  $a \times 2^3 = 8$

MODULE - III  
Sequences And  
Series

Notes

$$\therefore a = \frac{8}{8} = 1$$

Therefore, the G. P. is

$$1, 2, 4, 8, 16, \dots$$

**Example 13.17** Which term of the G. P.:  $5, -10, 20, -40, \dots$  is 320?

**Solution :** In this case,  $a = 5$ ;  $r = \frac{-10}{5} = -2$ .

Suppose that 320 is the  $n^{\text{th}}$  term of the G. P.

By the formula,  $t_n = ar^{n-1}$ , we get

$$t_n = 5 \cdot (-2)^{n-1}$$

$$\therefore 5 \cdot (-2)^{n-1} = 320 \quad \text{(Given)}$$

$$\therefore (-2)^{n-1} = 64 = (-2)^6$$

$$\therefore n - 1 = 6$$

$$\therefore n = 7$$

Hence, 320 is the 7<sup>th</sup> term of the G. P.

**Example 13.18** If  $a, b, c$ , and  $d$  are in G. P., then show that  $(a + b)^2$ ,  $(b + c)^2$ , and  $(c + d)^2$  are also in G. P.

**Solution.** Since  $a, b, c$ , and  $d$  are in G. P.,

$$\therefore \frac{b}{a} = \frac{c}{b} = \frac{d}{c}$$

$$\therefore b^2 = ac, c^2 = bd, ad = bc \quad \dots(1)$$

$$\text{Now, } (a + b)^2 (c + d)^2 = [(a + b)(c + d)]^2 = (ac + bc + ad + bd)^2$$

$$= (b^2 + c^2 + 2bc)^2 \quad \dots[\text{Using (1)}]$$

$$= [(b + c)^2]^2$$

$$\therefore \frac{(c + d)^2}{(b + c)^2} = \frac{(b + c)^2}{(a + b)^2}$$

Thus,  $(a + b)^2$ ,  $(b + c)^2$ ,  $(c + d)^2$  are in G. P.





Notes



### CHECK YOUR PROGRESS 13.4

1. The first term and the common ratio of a G. P. are respectively 3 and  $-\frac{1}{2}$ . Write down the first five terms.
2. Which term of the G. P. series 1, 2, 4, 8, 16, ... is 1024? Is 520 a term of the G. P. series.?
3. Three numbers are in G. P. Their sum is 43 and their product is 216. Find the numbers in proper order.
4. The  $n^{\text{th}}$  term of a G. P. is  $2 \times 3^n$  for all  $n$ . Find (a) the first term (b) the common ratio of the G. P.

### 13.6 SUM OF $n$ TERMS OF A G. P.

Let  $a$  denote the first term and  $r$  the common ratio of a G. P. Let  $S_n$  represent the sum of first  $n$  terms of the G. P.

$$\text{Thus, } S_n = a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} \quad \dots (1)$$

Multiplying (1) by  $r$ , we get

$$r S_n = ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} + ar^n \quad \dots (2)$$

$$(1) - (2) \Rightarrow S_n - rS_n = a - ar^n$$

$$\text{or } S_n (1 - r) = a (1 - r^n)$$

$$\therefore S_n = \frac{a(1 - r^n)}{1 - r} \quad \dots(A)$$

$$= \frac{a(r^n - 1)}{r - 1} \quad \dots(B)$$

Either (A) or (B) gives the sum up to the  $n^{\text{th}}$  term when  $r \neq 1$ . It is convenient to use formula (A) when  $|r| < 1$  and (B) when  $|r| > 1$ .

**Example 13.19** Find the sum of the G. P.:

1, 3, 9, 27, ... up to the 10<sup>th</sup> term.

**Solution :** Here the first term ( $a$ ) = 1 and the common ratio ( $r$ ) =  $\frac{3}{1} = 3$

Now using the formula,  $S_n = \frac{a(r^n - 1)}{r - 1}$ , ( $\because r > 1$ ) we get

$$S_{10} = \frac{1.(3^{10} - 1)}{3 - 1} = \frac{3^{10} - 1}{2}$$

MODULE - III  
Sequences And  
Series



Notes

**Example 13.20** Find the sum of the G. P.:  $\frac{1}{\sqrt{3}}, 1, \sqrt{3}, \dots, 81$

**Solution :** Here,  $a = \frac{1}{\sqrt{3}}$ ;  $r = \sqrt{3}$  and  $t_n = l = 81$

$$\text{Now } t_n = 81 = \frac{1}{\sqrt{3}} (\sqrt{3})^{n-1} = (\sqrt{3})^{n-2}$$

$$\therefore (\sqrt{3})^{n-2} = 3^4 = (\sqrt{3})^8$$

$$\therefore n - 2 = 8$$

$$\text{or } n = 10$$

$$\therefore S_n = \frac{\frac{1}{\sqrt{3}} [\sqrt{3}^{10} - 1]}{\sqrt{3} - 1}$$

$$= \frac{(\sqrt{3})^{10} - 1}{3 - \sqrt{3}}$$

**Example 13.21** Find the sum of the G. P.: 0.6, 0.06, 0.006, 0.0006, ... to  $n$  terms.

**Solution.** Here,  $a = 0.6 = \frac{6}{10}$  and  $r = \frac{0.06}{0.6} = \frac{1}{10}$

Using the formula  $S_n = \frac{a(1-r^n)}{1-r}$ , we have  $[\because r < 1]$

$$S_n = \frac{\frac{6}{10} \left[ 1 - \left( \frac{1}{10} \right)^n \right]}{1 - \frac{1}{10}} = \frac{6}{9} \left( 1 - \frac{1}{10^n} \right)$$

$$= \frac{2}{3} \left( 1 - \frac{1}{10^n} \right)$$

Hence, the required sum is  $\frac{2}{3} \left( 1 - \frac{1}{10^n} \right)$ .



**Example 13.22** How many terms of the following G.P.: 64, 32, 16, ... has the sum  $127\frac{1}{2}$  ?

**Solution :** Here,  $a = 64$ ,  $r = \frac{32}{64} = \frac{1}{2}$  ( $<1$ ) and  $S_n = 127\frac{1}{2} = \frac{255}{2}$ .

Using the formula  $S_n = \frac{a(1-r^n)}{1-r}$ , we get

$$S_n = \frac{64 \left\{ 1 - \left( \frac{1}{2} \right)^n \right\}}{1 - \frac{1}{2}}$$

$$\Rightarrow \frac{64 \left\{ 1 - \left( \frac{1}{2} \right)^n \right\}}{1 - \frac{1}{2}} = \frac{255}{2} \quad \dots \text{(given)}$$

$$\text{or } 128 \left[ 1 - \left( \frac{1}{2} \right)^n \right] = \frac{255}{2}$$

$$\text{or } 1 - \left( \frac{1}{2} \right)^n = \frac{255}{256}$$

$$\text{or } \left( \frac{1}{2} \right)^n = 1 - \frac{255}{256} = \frac{1}{256} = \left( \frac{1}{2} \right)^8$$

$$\therefore n = 8$$

Thus, the required number of terms is 8.

**Example 13.23** Find the sum of the following sequence :

2, 22, 222, ..... to  $n$  terms.

**Solution :** Let  $S$  denote the sum. Then

$$S = 2 + 22 + 222 + \dots \text{ to } n \text{ terms}$$

$$= 2 (1 + 11 + 111 + \dots \text{ to } n \text{ terms})$$

$$= \frac{2}{9} (9 + 99 + 999 + \dots \text{ to } n \text{ terms})$$

MODULE - III  
Sequences And  
Series



Notes

$$\begin{aligned}
 &= \frac{2}{9} \{ (10-1) + (10^2-1) + (10^3-1) + \dots \text{to } n \text{ terms} \} \\
 &= \frac{2}{9} \{ (10-10^2+10^3 + \dots \text{to } n \text{ terms}) - (1+1+1+ \dots \text{to } n \text{ terms}) \} \\
 &= \frac{2}{9} \left\{ \frac{(10^n-1)}{10-1} - n \right\} \quad [ \because 10-10^2+10^3 + \dots \text{ is a G P with } r = -10 < 1 ] \\
 &= \frac{2}{9} \left[ \frac{10^n-1-9n}{9} \right] \\
 &= \frac{2}{81} (10^n-1-9n)
 \end{aligned}$$

**Example 13.24** Find the sum up to  $n$  terms of the sequence:

0.7, 0.77, 0.777, ...

**Solution :** Let  $S$  denote the sum, then

$$\begin{aligned}
 S &= 0.7 + 0.77 + 0.777 + \dots \text{ to } n \text{ terms} \\
 &= 7(0.1 + 0.11 + 0.111 + \dots \text{ to } n \text{ terms}) \\
 &= \frac{7}{9} (0.9 + 0.99 + 0.999 + \dots \text{ to } n \text{ terms}) \\
 &= \frac{7}{9} \{ (1-0.1) + (1-0.01) + (1-0.001) + \dots \text{ to } n \text{ terms} \} \\
 &= \frac{7}{9} \{ (1+1+1+ \dots \text{ } n \text{ terms}) - (0.1 + 0.01 + 0.001 + \dots \text{ to } n \text{ terms}) \} \\
 &= \frac{7}{9} \left\{ n - \left( \frac{1}{10} + \frac{1}{10^2} + \frac{1}{10^3} + \dots \text{to } n \text{ terms} \right) \right\} \\
 &= \frac{7}{9} \left\{ n - \frac{\frac{1}{10} \left( 1 - \frac{1}{10^n} \right)}{1 - \frac{1}{10}} \right\} \quad (\text{Since } r < 1) \\
 &= \frac{7}{9} \left\{ n - \frac{1}{9} \left( 1 - \frac{1}{10^n} \right) \right\}
 \end{aligned}$$



$$= \frac{7}{9} \left[ \frac{9n - 1 + 10^{-n}}{9} \right]$$

$$= \frac{7}{81} [9n - 1 + 10^{-n}]$$



### CHECK YOUR PROGRESS 13.5

- Find the sum of each of the following G. P's :
  - 6, 12, 24, ... to 10 terms
  - $1, -\frac{1}{2}, \frac{1}{4}, -\frac{1}{8}, \frac{1}{16}, \dots$  to 20 terms.
- How many terms of the G. P. 8, 16, 32, 64, ... have their sum 8184 ?
- Show that the sum of the G. P.  $a + b + \dots + l$  is  $\frac{bl - a^2}{b - a}$
- Find the sum of each of the following sequences up to  $n$  terms.
  - 8, 88, 888, ...
  - 0.2, 0.22, 0.222, ...

### 13.7 INFINITE GEOMETRIC PROGRESSION

So far, we have found the sum of a finite number of terms of a G. P. We will now learn to find out the sum of infinitely many terms of a GP such as.

$$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$$

We will proceed as follows: Here  $a = 1, r = \frac{1}{2}$ .

The  $n^{\text{th}}$  term of the G. P. is  $t_n = \frac{1}{2^{n-1}}$  and sum to  $n$  terms

$$\text{i.e., } S_n = \frac{1 - \frac{1}{2^n}}{1 - \frac{1}{2}} = 2 \left( 1 - \frac{1}{2^n} \right)$$

$$= 2 - \frac{1}{2^{n-1}} < 2.$$

So, no matter, how large  $n$  may be, the sum of  $n$  terms is never more than 2.



So, if we take the sum of all the infinitely many terms, we shall not get more than 2 as answer.

Also note that the recurring decimal 0.3 is really  $0.3 + 0.03 + 0.003 + 0.0003 + \dots$

i.e., 0.3 is actually the sum of the above infinite sequence.

On the other hand it is at once obvious that if we sum infinitely many terms of the G. P. 1, 2, 4, 8, 16, ... we shall get a finite sum.

So, sometimes we may be able to add the infinitely many terms of G. P. and sometimes are may not. We shall discuss this question now.

### 13.7.1 SUM OF INFINITE TERMS OF A G. P.

Let us consider a G. P. with infinite number of terms and common ratio  $r$ .

**Case 1 :** We assume that  $|r| > 1$

The expression for the sum of  $n$  terms of the G. P. is then given by

$$\begin{aligned} S_n &= \frac{a(r^n - 1)}{r - 1} \\ &= \frac{ar^n}{r - 1} - \frac{a}{r - 1} \quad \dots (A) \end{aligned}$$

Now as  $n$  becomes larger and larger  $r^n$  also becomes larger and larger. Thus, when  $n$  is infinitely large and  $|r| > 1$  then the sum is also infinitely large which has no importance in Mathematics. We now consider the other possibility.

**Case 2 :** Let  $|r| < 1$

Formula (A) can be written as

$$S = \frac{a(1 - r^n)}{1 - r} = \frac{a}{1 - r} - \frac{ar^n}{1 - r}$$

Now as  $n$  becomes infinitely large,  $r^n$  becomes infinitely small, i.e., as  $n \rightarrow \infty$ ,  $r^n \rightarrow 0$ , then the above expression for sum takes the form

$$S = \frac{a}{1 - r}$$

Hence, the sum of an infinite G. P. with the first term  $a$  and common ratio  $r$  is given by

$$S = \frac{a}{1 - r}, \text{ when } |r| < 1 \quad \dots (i)$$



**Example 13.25** Find the sum of the infinite G.P.  $\frac{1}{3}, -\frac{2}{9}, \frac{4}{27}, -\frac{8}{81}, \dots$

**Solution :** Here, the first term of the infinite G.P. is  $a = \frac{1}{3}$ , and  $r = \frac{-\frac{2}{9}}{\frac{1}{3}} = -\frac{2}{3}$ .

$$\text{Here, } |r| = \left| -\frac{2}{3} \right| = \frac{2}{3} < 1$$

$\therefore$  Using the formula for sum  $S = \frac{a}{1-r}$  we have

$$S = \frac{\frac{1}{3}}{1 - \left(-\frac{2}{3}\right)} = \frac{\frac{1}{3}}{1 + \frac{2}{3}} = \frac{1}{5}$$

Hence, the sum of the given G.P. is  $\frac{1}{5}$ .

**Example 13.26** Express the recurring decimal  $0.\bar{3}$  as an infinite G.P. and find its value in rational form.

**Solution.**  $0.\bar{3} = 0.3333333 \dots$   
 $= 0.3 + 0.03 + 0.003 + 0.0003 + \dots$   
 $= \frac{3}{10} + \frac{3}{10^2} + \frac{3}{10^3} + \frac{3}{10^4} + \dots$

The above is an infinite G.P. with the first term  $a = \frac{3}{10}$  and  $r = \frac{\frac{3}{10^2}}{\frac{3}{10}} = \frac{1}{10} < 1$

Hence, by using the formula  $S = \frac{a}{1-r}$ , we get

$$0.\bar{3} = \frac{\frac{3}{10}}{1 - \frac{1}{10}} = \frac{\frac{3}{10}}{\frac{9}{10}}$$

$$= \frac{3}{9} = \frac{1}{3}$$

Hence, the recurring decimal  $0.\bar{3} = \frac{1}{3}$ .



**Example 13.27** The distance travelled (in cm) by a simple pendulum in consecutive seconds are 16, 12, 9, ... How much distance will it travel before coming to rest ?

**Solution :** The distance travelled by the pendulum in consecutive seconds are, 16, 12, 9, ... is an infinite geometric progression with the first term  $a = 16$  and  $r = \frac{12}{16} = \frac{3}{4} < 1$ .

Hence, using the formula  $S = \frac{a}{1-r}$  we have

$$S = \frac{16}{1 - \frac{3}{4}} = \frac{16}{\frac{1}{4}} = 64$$

$\therefore$  Distance travelled by the pendulum is 64 cm.

**Example 13.28** The sum of an infinite G.P. is 3 and sum of its first two terms is  $\frac{8}{3}$ . Find the first term.

**Solution:** In this problem  $S = 3$ . Let  $a$  be the first term and  $r$  be the common ratio of the given infinite G.P.

Then according to the question.

$$a + ar = \frac{8}{3}$$

$$\text{or, } 3a(1+r) = 8 \quad \dots (1)$$

Also from  $S = \frac{a}{1-r}$ , we have

$$3 = \frac{a}{1-r}$$

$$\text{or, } a = 3(1-r) \quad \dots (2)$$

From (1) and (2), we get.

$$3 \cdot 3(1-r)(1+r) = 8$$

$$\text{or, } 1 - r^2 = \frac{8}{9}$$





or,  $r^2 = \frac{1}{9}$

or,  $r = \pm \frac{1}{3}$

From (2),  $a = 3 \left( 1 \mp \frac{1}{3} \right) = 2$  or  $4$  according as  $r = \pm \frac{1}{3}$ .



**CHECK YOUR PROGRESS 13.6**

(1) Find the sum of each of the following infinite G. P.'s :

(a)  $1 + \frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \dots \infty$       (b)  $\frac{2}{5} + \frac{3}{5^2} + \frac{2}{5^3} + \frac{3}{5^4} + \dots \infty$

2. Express the following recurring decimals as an infinite G. P. and then find out their values as a rational number.

(a)  $0.\overline{7}$       (b)  $0.3\overline{15}$

3. The sum of an infinite G. P. is 15 and the sum of the squares of the terms is 45. Find the G.P.

4. The sum of an infinite G. P. is  $\frac{1}{3}$  and the first term is  $\frac{1}{4}$ . Find the G.P.

**13.8 GEOMETRIC MEAN (G. M.)**

If  $a, G, b$  are in G. P., then  $G$  is called the geometric mean between  $a$  and  $b$ .

If three numbers are in G. P., the middle one is called the geometric mean between the other two.

If  $a, G_1, G_2, \dots, G_n, b$  are in G. P.,

then  $G_1, G_2, \dots, G_n$  are called  $n$  G. M.'s between  $a$  and  $b$ .

The geometric mean of  $n$  numbers is defined as the  $n^{\text{th}}$  root of their product.

Thus if  $a_1, a_2, \dots, a_n$  are  $n$  numbers, then their

$$\text{G. M.} = (a_1, a_2, \dots, a_n)^{\frac{1}{n}}$$

Let  $G$  be the G. M. between  $a$  and  $b$ , then  $a, G, b$  are in G. P.

$$\frac{G}{a} = \frac{b}{G}$$

MODULE - III  
Sequences And  
Series



Notes

or,  $G^2 = ab$

or,  $G = \sqrt{ab}$

$\therefore$  Geometric mean  $= \sqrt{\text{Product of extremes}}$

Given any two positive numbers  $a$  and  $b$ , any number of geometric means can be inserted  
.....  $a_1, a_2, a_3, \dots, a_n$  be  $n$  geometric means between  $a$  and  $b$ .

Then  $a, a_1, a_2, \dots, a_n, b$  is a G. P.

Thus,  $b$  being the  $(n + 2)^{\text{th}}$  term, we have

$$b = a r^{n+1}$$

or,  $r^{n+1} = \frac{b}{a}$

or,  $r = \left(\frac{b}{a}\right)^{\frac{1}{n+1}}$

Hence,  $a_1 = ar = a \times \left(\frac{b}{a}\right)^{\frac{1}{n+1}}$

$$a_2 = ar^2 = a \times \left(\frac{b}{a}\right)^{\frac{2}{n+1}}$$

.....  
.....

$$a_n = ar^n = a \times \left(\frac{b}{a}\right)^{\frac{n}{n+1}}$$

Further we can show that the product of these  $n$  G. M.'s is equal to  $n^{\text{th}}$  power of the single geometric mean between  $a$  and  $b$ .

Multiplying  $a_1, a_2, \dots, a_n$ , we have

$$a_1, a_2 \cdots a_n = a^n \left(\frac{b}{a}\right)^{\frac{1}{n+1} + \frac{2}{n+1} + \cdots + \frac{n}{n+1}}$$

$$= a^n \left(\frac{b}{a}\right)^{\frac{1+2+\cdots+n}{n+1}}$$



$$\begin{aligned}
 &= a^n \left( \frac{b}{a} \right)^{\frac{n(n+1)}{2(n+1)}} \\
 &= a^n \left( \frac{b}{a} \right)^{\frac{n}{2}} \\
 &= (ab)^{\frac{n}{2}} = (\sqrt{ab})^n = G^n \\
 &= (\text{single G. M. between } a \text{ and } b)^n
 \end{aligned}$$

**Example 13.29** Find the G. M. between  $\frac{3}{2}$  and  $\frac{27}{2}$

**Solution :** We know that if  $a$  is the G. M. between  $a$  and  $b$ , then

$$G = \sqrt{ab}$$

$$\therefore \text{G. M. between } \frac{3}{2} \text{ and } \frac{27}{2} = \sqrt{\frac{3}{2} \times \frac{27}{2}} = \frac{9}{2}$$

**Example 13.30** Insert three geometric means between 1 and 256.

**Solution :** Let  $G_1, G_2, G_3$ , be the three geometric means between 1 and 256.

Then 1,  $G_1, G_2, G_3, 256$  are in G. P.

If  $r$  be the common ratio, then  $t_5 = 256$

$$\text{i.e., } ar^4 = 256 \Rightarrow 1 \cdot r^4 = 256$$

$$\text{or, } r^2 = 16$$

$$\text{or, } r = \pm 4$$

When  $r = 4$ ,  $G_1 = 1 \cdot 4 = 4$ ,  $G_2 = 1 \cdot (4)^2 = 16$  and  $G_3 = 1 \cdot (4)^3 = 64$

When  $r = -4$ ,  $G_1 = -4$ ,  $G_2 = (1) (-4)^2 = 16$  and  $G_3 = (1) (-4)^3 = -64$

$\therefore$  G.M. between 1 and 256 are 4, 16, 64, or,  $-4, 16, -64$ .

**Example 13.31** If 4, 36, 324 are in G. P. insert two more numbers in this progression so that it again forms a G. P.

**Solution :** G. M. between 4 and 36 =  $\sqrt{4 \times 36} = \sqrt{144} = 12$

$$\text{G. M. between 36 and } 324 = \sqrt{36 \times 324} = 6 \times 18 = 108$$

MODULE - III  
Sequences And  
Series



Notes

If we introduce 12 between 4 and 36 and 108 between 36 and 324, the numbers 4, 12, 36, 108, 324 form a G. P.

∴ The two new numbers inserted are 12 and 108.

**Example 13.32** Find the value of  $n$  such that  $\frac{a^{n+1} + b^{n+1}}{a^n + b^n}$  may be the geometric mean between  $a$  and  $b$ .

**Solution :** If  $x$  be G. M. between  $a$  and  $b$ , then

$$x = a^{\frac{1}{2}} \times b^{\frac{1}{2}}$$

$$\therefore \frac{a^{n+1} + b^{n+1}}{a^n + b^n} = a^{\frac{1}{2}} b^{\frac{1}{2}}$$

$$\text{or, } a^{n+1} + b^{n+1} = \left( a^{\frac{1}{2}} b^{\frac{1}{2}} \right) (a^n + b^n)$$

$$\text{or, } a^{n+1} + b^{n+1} = a^{n+\frac{1}{2}} b^{\frac{1}{2}} + a^{\frac{1}{2}} b^{n+\frac{1}{2}}$$

$$\text{or, } a^{n+1} - a^{n+\frac{1}{2}} \cdot b^{\frac{1}{2}} = a^{\frac{1}{2}} b^{n+\frac{1}{2}} - b^{n+1}$$

$$\text{or, } a^{n+\frac{1}{2}} \left( a^{\frac{1}{2}} - b^{\frac{1}{2}} \right) = b^{n+\frac{1}{2}} \left( a^{\frac{1}{2}} - b^{\frac{1}{2}} \right)$$

$$\text{or, } a^{n+\frac{1}{2}} = b^{n+\frac{1}{2}}$$

$$\text{or, } \frac{a^{n+\frac{1}{2}}}{b^{n+\frac{1}{2}}} = 1$$

$$\text{or, } \left( \frac{a}{b} \right)^{n+\frac{1}{2}} = \left( \frac{a}{b} \right)^0$$

$$\therefore n + \frac{1}{2} = 0 \text{ or, } n = -\frac{1}{2}$$



### 13.8.1 RELATIONSHIP BETWEEN A. M. AND G.M.

Let  $a$  and  $b$  be the two numbers.

Let  $A$  and  $G$  be the A. M. and G. M. respectively between  $a$  and  $b$

$$\therefore A = \frac{a+b}{2}, G = \sqrt{ab}$$

$$\begin{aligned} A - G &= \frac{a+b}{2} - \sqrt{ab} \\ &= \frac{(\sqrt{a})^2 + (\sqrt{b})^2 - 2\sqrt{ab}}{2} \\ &= \frac{1}{2}(\sqrt{a} - \sqrt{b})^2 > 0 \end{aligned}$$

$$\therefore A > G$$

**Example 13.33** The arithmetic mean between two numbers is 34 and their geometric mean is 16. Find the numbers.

**Solution :** Let the numbers be  $a$  and  $b$ .

Since A. M. between  $a$  and  $b$  is 34,

$$\therefore \frac{a+b}{2} = 34, \text{ or, } a+b = 68 \quad \dots (1)$$

Since G. M. between  $a$  and  $b$  is 16,

$$\therefore \sqrt{ab} = 16 \text{ or, } ab = 256$$

$$\text{we know that } (a-b)^2 = (a+b)^2 - 4ab \quad \dots (2)$$

$$\begin{aligned} &= (68)^2 - 4 \times 256 \\ &= 4624 - 1024 = 3600 \end{aligned}$$

$$\therefore a-b = \sqrt{3600} = 60 \quad \dots (3)$$

Adding (1) and (3), we get,  $2a = 128$

$$\therefore a = 64$$

Subtracting (3) from (1), we get

$$2b = 8 \quad \text{or, } b = 4$$

$\therefore$  Required numbers are 64 and 4.



Notes

**Example 13.34** The arithmetic mean between two quantities  $b$  and  $c$  is  $a$  and the two geometric means between them are  $g_1$  and  $g_2$ . Prove that

$$g_1^3 + g_2^3 = 2abc$$

**Solution :** The A. M. between  $b$  and  $c$  is  $a$

$$\therefore \frac{b+c}{2} = a, \text{ or, } b+c = 2a$$

Again  $g_1$  and  $g_2$  are two G. M.'s between  $b$  and  $c$

$$\therefore b, g_1, g_2, c \text{ are in G. P.}$$

If  $r$  be the common ratio, then

$$c = br^3 \quad \text{or,} \quad r = \left(\frac{c}{b}\right)^{\frac{1}{3}}$$

$$g_1 = br = b\left(\frac{c}{b}\right)^{\frac{1}{3}}$$

$$\text{and } g_2 = br^2 = b\left(\frac{c}{b}\right)^{\frac{2}{3}}$$

$$\begin{aligned} \therefore g_1^3 + g_2^3 &= b^3 \left[ \left(\frac{c}{b}\right) + \left(\frac{c}{b}\right)^2 \right] \\ &= b^3 \times \frac{c}{b} \left(1 + \frac{c}{b}\right) \\ &= b^2 c \times \left(\frac{b+c}{b}\right) \\ &= bc(2a) \quad [\text{since } b+c = 2a] \\ &= 2abc \end{aligned}$$

**Example 13.35** If one geometric mean  $G$  and two arithmetic means  $p$  and  $q$  be inserted between two quantities, show that :

$$G^2 = (2p - q)(2q - p)$$

**Solution :** Let the two quantities be  $a$  and  $b$ , then

$$G = \sqrt{ab} \quad \text{or, } G^2 = ab \quad \dots(i)$$



Also  $p$  and  $q$  are two A. M.'s between  $a$  and  $b$

$$\therefore a, p, q, b \text{ are in A. P.}$$

$$\therefore p - a = q - p \text{ and } q - p = b - q$$

$$\therefore a = 2p - q \text{ and } b = 2q - p$$

$$\therefore G^2 = ab = (2p - q)(2q - p)$$

**Example 13.36** The product of first three terms of a G. P. is 1000. If we add 6 to its second term and 7 to its 3rd term, the three terms form an A. P. Find the terms of the G. P.

**Solution :** Let  $t_1 = \frac{a}{r}, t_2 = a$  and  $t_3 = ar$  be the first three terms of G. P.

Then, their product =  $\frac{a}{r} \cdot a \cdot ar = 1000$  or,  $a^3 = 1000$ , or,  $a = 10$

By the question,  $t_1, t_2 + 6, t_3 + 7$  are in A. P. ... (1)

i.e.  $\frac{a}{r}, a + 6, ar + 7$  are in A. P.

$$\therefore (a + 6) - \frac{a}{r} = (ar + 7) - (a + 6)$$

$$\text{or, } 2(a + 6) = \frac{a}{r} + (ar + 7)$$

$$\text{or, } 2(10 + 6) = \frac{10}{r} + (10r + 7) \quad [\text{using (1)}]$$

$$\text{or, } 32r = 10 + 10r^2 + 7r$$

$$\text{or, } 10r^2 - 25r + 10 = 0$$

$$\therefore r = \frac{25 \pm \sqrt{625 - 400}}{20} = \frac{25 \pm 15}{20}$$

$$= 2, \frac{1}{2}$$

When  $a = 10, r = 2$ . then the terms are  $\frac{10}{2}, 10(2)$  i.e., 5, 10, 20

When  $a = 10, r = \frac{1}{2}$  then the terms are  $10(2), 10, 10\left(\frac{1}{2}\right)$  i.e., 20, 10, 5



Notes

**CHECK YOUR PROGRESS 13.7**

1. Insert 8 G. M.'s between 8 and  $\frac{1}{64}$ .
2. If  $a_1$  is the first of  $n$  geometric means between  $a$  and  $b$ , show that  $a_1^{n+1} = a^n b$
3. If  $G$  is the G. M. between  $a$  and  $b$ , prove that  $\frac{1}{G^2 - a^2} + \frac{1}{G^2 - b^2} = \frac{1}{G^2}$
4. If the A. M. and G. M. between two numbers are in the ratio  $m : n$ , then prove that the numbers are in the ratio  $m + \sqrt{m^2 - n^2} : m - \sqrt{m^2 - n^2}$
5. If  $A$  and  $G$  are respectively arithmetic and geometric means between two numbers  $a$  and  $b$ , then show that  $A > G$ .
6. The sum of first three terms of a G. P. is  $\frac{13}{12}$  and their product is  $-1$ . Find the G. P.
7. The product of three terms of a G. P. is 512. If 8 added to first and 6 added to second term, the numbers form an A. P., Find the numbers.

**LET US SUM UP**

- The general term of an A. P.  
 $a, a + d, a + 2d, \dots$  is given by  $t_n = a + (n - 1)d$
- $S_n$ , the sum of the first  $n$  terms of an A. P.  
 $a, a + d, a + 2d, \dots$  is  
given by  $S_n = \frac{n}{2} [2a + (n - 1)d]$   
 $= \frac{n}{2} (a + l)$ , where  $l = a + (n - 1)d$ .
- A sequence in which the difference of two consecutive terms is always constant ( $\neq 0$ ) is called an Arithmetic Progression (A. P.)
- $t_n = S_n - S_{n-1}$





- $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$
- An arithmetic mean between  $a$  and  $b$  is  $\frac{a+b}{2}$ .
- A sequence in which the ratio of two consecutive terms is always constant ( $\neq 0$ ) is called a Geometric Progression (G. P.)
- The  $n^{\text{th}}$  term of a G. P.:  $a, ar, ar^2, \dots$  is  $ar^{n-1}$
- Sum of the first  $n$  terms of a G. P.:  $a, ar, ar^2, \dots$  is

$$S_n = \frac{a(r^n - 1)}{r - 1} \text{ for } |r| > 1$$

$$= \frac{a(1 - r^n)}{1 - r} \text{ for } |r| < 1$$

- The sums of an infinite G. P.  $a, ar, ar^2, \dots$  is given by

$$S = \frac{a}{1 - r} \text{ for } |r| < 1$$

- Geometric mean  $G$  between two numbers  $a$  and  $b$  is  $\sqrt{ab}$
- The arithmetic mean  $A$  between two numbers  $a$  and  $b$  is always greater than the corresponding Geometric mean  $G$  i.e.,  $A > G$ .



### SUPPORTIVE WEB SITES

<http://www.wikipedia.org>

<http://mathworld.wolfram.com>



### TERMINAL EXERCISE

1. Find the sum of all the natural numbers between 100 and 200 which are divisible by 7.
2. The sum of the first  $n$  terms of two A. P.'s are in the ratio  $(2n - 1) : (2n + 1)$ . Find the ratio of their 10<sup>th</sup> terms.
3. If  $a, b, c$  are in A. P. then show that  $b + c, c + a, a + b$  are also in A. P.

MODULE - III  
Sequences And  
Series



## Notes

4. If  $a_1, a_2, \dots, a_n$  are in A. P., then prove that

$$\frac{1}{a_1 a_2} + \frac{1}{a_2 a_3} + \frac{1}{a_3 a_4} + \dots + \frac{1}{a_{n-1} a_n} = \frac{n-1}{a_1 a_n}$$

5. If  $(b-c)^2, (c-a)^2, (a-b)^2$  are in A. P., then prove that

$$\frac{1}{b-c}, \frac{1}{c-a}, \frac{1}{a-b}, \text{ are also in A. P.}$$

6. If the  $p^{\text{th}}, q^{\text{th}}$  and  $r^{\text{th}}$  terms are  $P, Q, R$  respectively. Prove that  $P(Q-R) + Q(R-P) + r(P-Q) = 0$ .

7. If  $a, b, c$  are in G. P. then prove that

$$a^2 b^2 c^2 \left( \frac{1}{a^3} + \frac{1}{b^3} + \frac{1}{c^3} \right) = a^3 + b^3 + c^3$$

8. If  $a, b, c, d$  are in G. P., show that each of the following form a G. P. :

(a)  $(a^2 - b^2), (b^2 - c^2), (c^2 - d^2)$

(b)  $\frac{1}{a^2 + b^2}, \frac{1}{b^2 + c^2}, \frac{1}{c^2 + d^2}$

9. If  $x, y, z$  are the  $p^{\text{th}}, q^{\text{th}}$  and  $r^{\text{th}}$  terms of a G. P., prove that

$$x^{q-r} y^{r-p} z^{p-q} = 1$$

10. If  $a, b, c$  are in A. P. and  $x, y, z$  are in G. P. then prove that

$$x^{b-c} y^{c-a} z^{a-b} = 1$$

11. If the sum of the first  $n$  terms of a G. P. is represented by  $S_n$ , then prove that

$$S_n (S_{3n} - S_{2n}) = (S_{2n} - S_n)^2$$

12. If  $p, q, r$  are in A. P. then prove that the  $p^{\text{th}}, q^{\text{th}}$  and  $r^{\text{th}}$  terms of a G. P. are also in G. P.

13. If  $S_n = 1 + \frac{1}{2} + \frac{1}{2^2} \dots + \frac{1}{2^{n-1}}$ , find the least value of  $n$  such that

$$2 - S_n < \frac{1}{100}$$

14. If the sum of the first  $n$  terms of a G. P. is  $S$  and the product of these terms is  $p$  and the sum of their reciprocals is  $R$ , then prove that

$$p^2 = \left( \frac{S}{R} \right)^n$$



**ANSWERS**



Notes

**CHECK YOUR PROGRESS 13.1**

1. (a)  $2n - 1$       (b)  $2n + 1$   
 2. 3, 5, 7, 9, ...      3. 20, 16      4. no      5.  $m + n$       6. 10, 6, 2, ...

**CHECK YOUR PROGRESS 13.2**

1. (a) 435      (b)  $\frac{n}{2}[21 - 5n^2]$   
 2. 5      3. 12  
 4. 37      5.  $n^2 + 9n, 9$   
 6.  $2a$

**CHECK YOUR PROGRESS 13.3**

2. 5

**CHECK YOUR PROGRESS 13.4**

1.  $3, -\frac{3}{2}, \frac{3}{4}, -\frac{3}{8}, \frac{3}{16}$       2. 11<sup>th</sup>, no  
 3. 36, 6, 1 or 1, 6, 36      4. (a) 6      (b) 3

**CHECK YOUR PROGRESS 13.5**

1. (a) 6138      (b)  $\frac{2}{3}\left(1 - \frac{1}{2^{20}}\right)$   
 2. 10.  
 4. (a)  $\frac{80}{81}(10^n - 1) - \frac{8n}{9}$       (b)  $\frac{2n}{9} - \frac{2}{81}\left(1 - \frac{1}{10^n}\right)$

MODULE - III  
Sequences And  
Series



Notes

**CHECK YOUR PROGRESS 13.6**

1. (a)  $\frac{3}{2}$                       (b)  $\frac{13}{24}$

2. (a)  $\frac{7}{9}$                         (b)  $\frac{52}{165}$

3.  $5, \frac{10}{3}, \frac{20}{9}, \frac{40}{27}, \dots \infty$

4.  $\frac{1}{4}, \frac{1}{4^2}, \frac{1}{4^3}, \frac{1}{4^4}, \dots \infty$

**CHECK YOUR PROGRESS 13.7**

1.  $4, 2, 1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \dots$

6.  $\frac{4}{3}, -1, \frac{3}{4}$  or  $\frac{3}{4}, -1, \frac{4}{3}$

7.  $4, 8, 16$

**TERMINAL EXERCISE**

1. 2107

2. 37 : 39