



25

MAXIMA AND MINIMA

You are aware that in any transaction the total amount paid increases with the number of items purchased. Consider a function as $f(x) = 2x + 1, x > 0$. Let the function $f(x)$ represent the amount required for purchasing 'x' number of items.

The graph of the function $y = f(x) = 2x + 1, x > 0$ for different values of x is shown in Fig. 25.1.




Fig. 25.1

A look at the graph of the above function prompts us to believe that the function is increasing for positive values of x i.e., for $x > 0$. Can you think of another example in which value of the function decreases when x increases? Any such relation would be relationship between time and manpower/person(s) involved. You have learnt that they are inversely proportional. In other words we can say that time required to complete a certain work increases when number of persons (manpower) involved decreases and vice-versa. Consider such similar function as

$$g(x) = \frac{2}{x}, x > 0$$

The values of the function $g(x)$ for different values of x are plotted in Fig. 25.2.

MODULE - V
Calculus


Notes




Fig. 25.2

All these examples, despite the diversity of the variables involved, have one thing in common : function is either increasing or decreasing.

In this lesson, we will discuss such functions and their characteristics. We will also find out intervals in which a given function is increasing/decreasing and its application to problems on maxima and minima.


OBJECTIVES

After studying this lesson, you will be able to :

- define monotonic (increasing and decreasing) functions;
- establish that $\frac{dy}{dx} > 0$ in an interval for an increasing function and $\frac{dy}{dx} < 0$ for a decreasing function;
- define the points of maximum and minimum values as well as local maxima and local minima of a function from the graph;
- establish the working rule for finding the maxima and minima of a function using the first and the second derivatives of the function; and
- work out simple problems on maxima and minima.

EXPECTED BACKGROUND KNOWLEDGE

- Concept of function and types of function
- Differentiation of a function
- Solutions of equations and the inequations

25.1 INCREASING AND DECREASING FUNCTIONS

You have already seen the common trends of an increasing or a decreasing function. Here we will try to establish the condition for a function to be an increasing or a decreasing.

Maxima and Minima

Let a function $f(x)$ be defined over the closed interval $[a, b]$.

Let $x_1, x_2 \in [a, b]$, then the function $f(x)$ is said to be an increasing function in the given interval if $f(x_2) \geq f(x_1)$ whenever $x_2 > x_1$. It is said to be strictly increasing if $f(x_2) > f(x_1)$ for all $x_2 > x_1, x_1, x_2 \in [a, b]$.

In Fig. 25.3, $\sin x$ increases from -1 to $+1$ as x increases from $-\frac{\pi}{2}$ to $+\frac{\pi}{2}$.




Fig. 25.3

Note : A function is said to be an increasing function in an interval if $f(x+h) > f(x)$ for all x belonging to the interval when h is positive.

A function $f(x)$ defined over the closed interval $[a, b]$ is said to be a decreasing function in the given interval, if $f(x_2) \leq f(x_1)$, whenever $x_2 > x_1, x_1, x_2 \in [a, b]$. It is said to be strictly decreasing if $f(x_1) > f(x_2)$ for all $x_2 > x_1, x_1, x_2 \in [a, b]$.

In Fig. 25.4, $\cos x$ decreases from 1 to -1 as x increases from 0 to π .





Fig. 25.4



MODULE - V
Calculus

Notes

Note : A function is said to be a decreasing in an interval if $f(x+h) < f(x)$ for all x belonging to the interval when h is positive.

25.2 MONOTONIC FUNCTIONS

Let x_1, x_2 be any two points such that $x_1 < x_2$ in the interval of definition of a function $f(x)$. Then a function $f(x)$ is said to be monotonic if it is either increasing or decreasing. It is said to be monotonically increasing if $f(x_2) \geq f(x_1)$ for all $x_2 > x_1$ belonging to the interval and monotonically decreasing if $f(x_1) \geq f(x_2)$.

Example 25.1 Prove that the function $f(x) = 4x + 7$ is monotonic for all values of $x \in \mathbb{R}$.

Solution : Consider two values of x say $x_1, x_2 \in \mathbb{R}$

such that $x_2 > x_1$ (1)

Multiplying both sides of (1) by 4, we have $4x_2 > 4x_1$ (2)

Adding 7 to both sides of (2), to get

$$4x_2 + 7 > 4x_1 + 7$$

We have $f(x_2) > f(x_1)$

Thus, we find $f(x_2) > f(x_1)$ whenever $x_2 > x_1$.

Hence the given function $f(x) = 4x + 7$ is monotonic function. (monotonically increasing).

Example 25.2 Show that

$$f(x) = x^2$$

is a strictly decreasing function for all $x < 0$.

Solution : Consider any two values of x say x_1, x_2 such that

$$x_2 > x_1, \quad x_1, x_2 < 0 \quad \text{.....(i)}$$

Order of the inequality reverses when it is multiplied by a negative number. Now multiplying (i) by x_2 , we have

$$x_2 \cdot x_2 < x_1 \cdot x_2$$

or, $x_2^2 < x_1 x_2$ (ii)

Now multiplying (i) by x_1 , we have

$$x_1 \cdot x_2 < x_1 \cdot x_1$$

or, $x_1 x_2 < x_1^2$ (iii)

From (ii) and (iii), we have

$$x_2^2 < x_1 x_2 < x_1^2$$



or, $x_2^2 < x_1^2$
 or, $f(x_2) < f(x_1)$ (iv)

Thus, from (i) and (iv), we have for

$$x_2 > x_1,$$

$$f(x_2) < f(x_1)$$

Hence, the given function is strictly decreasing for all $x < 0$.



CHECK YOUR PROGRESS 25.1

1. (a) Prove that the function

$$f(x) = 3x + 4$$

is monotonic increasing function for all values of $x \in \mathbb{R}$.

- (b) Show that the function

$$f(x) = 7 - 2x$$

is monotonically decreasing function for all values of $x \in \mathbb{R}$.

- (c) Prove that $f(x) = ax + b$ where a, b are constants and $a > 0$ is a strictly increasing function for all real values of x .

2. (a) Show that $f(x) = x^2$ is a strictly increasing function for all real $x > 0$.

- (b) Prove that the function $f(x) = x^2 - 4$ is monotonically increasing for $x > 2$ and monotonically decreasing for $-2 < x < 2$ where $x \in \mathbb{R}$.

Theorem 1 : If $f(x)$ is an increasing function on an open interval $]a, b[$, then its derivative $f'(x)$ is positive at this point for all $x \in]a, b[$.

Proof : Let (x, y) or $[x, f(x)]$ be a point on the curve $y = f(x)$

For a positive δx , we have

$$x + \delta x > x$$

Now, function $f(x)$ is an increasing function

$$\therefore f(x + \delta x) > f(x)$$

or, $f(x + \delta x) - f(x) > 0$

or, $\frac{f(x + \delta x) - f(x)}{\delta x} > 0$ [$\because \delta x > 0$]

Taking δx as a small positive number and proceeding to limit, when $\delta x \rightarrow 0$

$$\lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x} > 0$$

MODULE - V
Calculus



Notes

or, $f'(x) > 0$

Thus, if $y = f(x)$ is an increasing function at a point, then $f'(x)$ is positive at that point.

Theorem 2 : If $f(x)$ is a decreasing function on an open interval $]a, b[$ then its derivative $f'(x)$ is negative at that point for all $x \in]a, b[$.

Proof : Let (x, y) or $[x, f(x)]$ be a point on the curve $y = f(x)$

For a positive δx , we have $x + \delta x > x$

Since the function is a decreasing function

$\therefore f(x + \delta x) < f(x) \qquad \delta x > 0$

or, $f(x + \delta x) - f(x) < 0$

Dividing by δx , we have $\frac{f(x + \delta x) - f(x)}{\delta x} < 0 \quad \delta x > 0$

or, $\lim_{\delta x \rightarrow 0} \frac{f(x + \delta x) - f(x)}{\delta x} < 0$

or, $f'(x) < 0$

Thus, if $y = f(x)$ is a decreasing function at a point, then, $f'(x)$ is negative at that point.

Note : If $f(x)$ is derivable in the closed interval $[a, b]$, then $f(x)$ is

- (i) increasing over $[a, b]$, if $f'(x) > 0$ in the open interval $]a, b[$
- (ii) decreasing over $[a, b]$, if $f'(x) < 0$ in the open interval $]a, b[$.

25.3 RELATION BETWEEN THE SIGN OF THE DERIVATIVE AND MONOTONICITY OF FUNCTION

Consider a function whose curve is shown in the Fig. 25.5




Fig. 25.5

We divide, our study of relation between sign of derivative of a function and its increasing or



Notes

decreasing nature (monotonicity) into various parts as per Fig. 25.5

- (i) P to R (ii) R to T (iii) T to V

(i) We observe that the ordinate (y-coordinate) for every succeeding point of the curve from P to R increases as also its x-coordinate. If (x_2, y_2) are the coordinates of a point that succeeds (x_1, y_1) then $x_2 > x_1$ yields $y_2 > y_1$ or $f(x_2) > f(x_1)$.

Also the tangent at every point of the curve between P and R makes acute angle with the positive direction of x-axis and thus the slope of the tangent at such points of the curve (except at R) is positive. At R where the ordinate is maximum the tangent is parallel to x-axis, as a result the slope of the tangent at R is zero.

We conclude for this part of the curve that

- (a) The function is monotonically increasing from P to R.
- (b) The tangent at every point (except at R) makes an acute angle with positive direction of x-axis.
- (c) The slope of tangent is positive i.e. $\frac{dy}{dx} > 0$ for all points of the curve for which y is increasing.
- (d) The slope of tangent at R is zero i.e. $\frac{dy}{dx} = 0$ where y is maximum.

(ii) The ordinate for every point between R and T of the curve decreases though its x-coordinate increases. Thus, for any point $x_2 > x_1$ yields $y_2 < y_1$, or $f(x_2) < f(x_1)$.

Also the tangent at every point succeeding R along the curve makes obtuse angle with positive direction of x-axis. Consequently, the slope of the tangent is negative for all such points whose ordinate is decreasing. At T the ordinate attains minimum value and the tangent is parallel to x-axis and as a result the slope of the tangent at T is zero.

We now conclude :

- (a) The function is monotonically decreasing from R to T.
- (b) The tangent at every point, except at T, makes obtuse angle with positive direction of x-axis.
- (c) The slope of the tangent is negative i.e., $\frac{dy}{dx} < 0$ for all points of the curve for which y is decreasing.

(d) The slope of the tangent at T is zero i.e. $\frac{dy}{dx} = 0$ where the ordinate is minimum.

(iii) Again, for every point from T to V

The ordinate is constantly increasing, the tangent at every point of the curve between T and V makes acute angle with positive direction of x-axis. As a result of which the slope of the tangent at each of such points of the curve is positive.

Conclusively,

$$\frac{dy}{dx} > 0$$

MODULE - V
Calculus

Notes

at all such points of the curve except at T and V, where $\frac{dy}{dx} = 0$. The derivative $\frac{dy}{dx} < 0$ on one side, $\frac{dy}{dx} > 0$ on the other side of points R, T and V of the curve where $\frac{dy}{dx} = 0$.

Example 25.3 Find for what values of x , the function

$$f(x) = x^2 - 6x + 8$$

is increasing and for what values of x it is decreasing.

Solution : $f(x) = x^2 - 6x + 8$

$$f'(x) = 2x - 6$$

For $f(x)$ to be increasing, $f'(x) > 0$

i.e., $2x - 6 > 0$ or, $2(x - 3) > 0$

or, $x - 3 > 0$ or, $x > 3$

The function increases for $x > 3$.

For $f(x)$ to be decreasing,

$$f'(x) < 0$$

i.e., $2x - 6 < 0$ or, $x - 3 < 0$

or, $x < 3$

Thus, the function decreases for $x < 3$.

Example 25.4 Find the interval in which $f(x) = 2x^3 - 3x^2 - 12x + 6$ is increasing or decreasing.

Solution : $f(x) = 2x^3 - 3x^2 - 12x + 6$

$$f'(x) = 6x^2 - 6x - 12$$

$$= 6(x^2 - x - 2)$$

$$= 6(x - 2)(x + 1)$$

For $f(x)$ to be increasing function of x ,

$$f'(x) > 0$$

i.e. $6(x - 2)(x + 1) > 0$ or, $(x - 2)(x + 1) > 0$

Since the product of two factors is positive, this implies either both are positive or both are negative.

Either $x - 2 > 0$ and $x + 1 > 0$ or $x - 2 < 0$ and $x + 1 < 0$

i.e. $x > 2$ and $x > -1$
 $x > 2$ implies $x > -1$
 $\therefore x > 2$

i.e. $x < 2$ and $x < -1$
 $x < -1$ implies $x < 2$
 $\therefore x < -1$

Hence $f(x)$ is increasing for $x > 2$ or $x < -1$.

Now, for $f(x)$ to be decreasing,

$$f(x) < 0$$

or, $6(x-2)(x+1) < 0$ or, $(x-2)(x+1) < 0$

Since the product of two factors is negative, only one of them can be negative, the other positive.

Therefore,

Either

$$x - 2 > 0 \text{ and } x + 1 < 0$$

i.e. $x > 2$ and $x < -1$

There is no such possibility
 that $x > 2$ and at the same time
 $x < -1$

or

$$x - 2 < 0 \text{ and } x + 1 > 0$$

i.e. $x < 2$ and $x > -1$

This can be put in this form

$$-1 < x < 2$$

\therefore The function is decreasing in $-1 < x < 2$.

Example 25.5 Determine the intervals for which the function

$$f(x) = \frac{x}{x^2 + 1} \text{ is increasing or decreasing.}$$

Solution : $f'(x) = \frac{(x^2 + 1) \frac{dx}{dx} - x \frac{d}{dx}(x^2 + 1)}{(x^2 + 1)^2}$

$$= \frac{(x^2 + 1) - x \cdot (2x)}{(x^2 + 1)^2}$$

$$= \frac{1 - x^2}{(x^2 + 1)^2}$$

$\therefore f'_x = \frac{(1-x)(1+x)}{(x^2 + 1)^2}$

As $(x^2 + 1)^2$ is positive for all real x .

Therefore, if $-1 < x < 0$, $(1 - x)$ is positive and $(1 + x)$ is positive, so $f'(x) > 0$;

\therefore If $0 < x < 1$, $(1 - x)$ is positive and $(1 + x)$ is positive, so $f'(x) > 0$;



MODULE - V
Calculus



Notes

If $x < -1$, $(1 - x)$ is positive and $(1 + x)$ is negative, so $f'(x) < 0$;

$x > 1$, $(1 - x)$ is negative and $(1 + x)$ is positive, so $f'(x) < 0$;

Thus we conclude that

the function is increasing for $-1 < x < 0$ and $0 < x < 1$

or, for $-1 < x < 1$

and the function is decreasing for $x < -1$ or $x > 1$

Note : Points where $f'(x) = 0$ are critical points. Here, critical points are $x = -1$, $x = 1$.

Example 25.6 Show that

(a) $f(x) = \cos x$ is decreasing in the interval $0 \leq x \leq \pi$.

(b) $f(x) = x - \cos x$ is increasing for all x .

Solution :(a) $f(x) = \cos x$

$$f'(x) = -\sin x$$

$f(x)$ is decreasing

If $f'(x) < 0$

i.e., $-\sin x < 0$

i.e., $\sin x > 0$

$\sin x$ is positive in the first quadrant and in the second quadrant, therefore, $\sin x$ is positive in $0 \leq x \leq \pi$

$\therefore f(x)$ is decreasing in $0 \leq x \leq \pi$

(b) $f(x) = x - \cos x$

$$f'(x) = 1 - (-\sin x)$$

$$= 1 + \sin x$$

Now, we know that the minimum value of $\sin x$ is -1 and its maximum; value is 1 i.e., $\sin x$ lies between -1 and 1 for all x ,

i.e., $-1 \leq \sin x \leq 1$ or $1 - 1 \leq 1 + \sin x \leq 1 + 1$

or $0 \leq 1 + \sin x \leq 2$

or $0 \leq f'(x) \leq 2$

or $0 \leq f'(x)$

or $f'(x) \geq 0$

$\Rightarrow f(x) = x - \cos x$ is increasing for all x .



CHECK YOUR PROGRESS 25.2

Find the intervals for which the following functions are increasing or decreasing.

1. (a) $f(x) = x^2 - 7x + 10$ (b) $f(x) = 3x^2 - 15x + 10$
2. (a) $f(x) = x^3 - 6x^2 - 36x + 7$ (b) $f(x) = x^3 - 9x^2 + 24x + 12$
3. (a) $y = -3x^2 - 12x + 8$ (b) $f(x) = 1 - 12x - 9x^2 - 2x^3$
4. (a) $y = \frac{x-2}{x+1}, x \neq -1$ (b) $y = \frac{x^2}{x-1}, x \neq 1$ (c) $y = \frac{x}{2} + \frac{2}{x}, x \neq 0$
5. (a) Prove that the function $\log \sin x$ is decreasing in $\left[\frac{\pi}{2}, \pi\right]$
 (b) Prove that the function $\cos x$ is increasing in the interval $[\pi, 2\pi]$
 (c) Find the intervals in which the function $\cos\left(2x + \frac{\pi}{4}\right), 0 \leq x \leq \pi$ is decreasing or increasing.

Find also the points on the graph of the function at which the tangents are parallel to x-axis.

25.4 MAXIMUM AND MINIMUM VALUES OF A FUNCTION

We have seen the graph of a continuous function. It increases and decreases alternatively. If the value of a continuous function increases upto a certain point then begins to decrease, then this point is called point of maximum and corresponding value at that point is called maximum value of the function. A stage comes when it again changes from decreasing to increasing. If the value of a continuous function decreases to a certain point and then begins to increase, then value at that point is called minimum value of the function and the point is called point of minimum.




Fig. 25.6

Fig. 25.6 shows that a function may have more than one maximum or minimum values. So, for continuous function we have maximum (minimum) value in an interval and these values are not absolute maximum (minimum) of the function. For this reason, we sometimes call them as local maxima or local minima.



Notes

MODULE - V
Calculus



Notes

A function $f(x)$ is said to have a maximum or a local maximum at the point $x = a$ where $a - b < a < a + b$ (See Fig. 25.7), if $f(a) \geq f(a \pm b)$ for all sufficiently small positive b .




Fig. 25.7




Fig. 25.8

A maximum (or local maximum) value of a function is the one which is greater than all other values on either side of the point in the immediate neighbourhood of the point.

A function $f(x)$ is said to have a minimum (or local minimum) at the point $x = a$ if $f(a) \leq f(a \pm b)$ where $a - b < a < a + b$

for all sufficiently small positive b .

In Fig. 25.8, the function $f(x)$ has local minimum at the point $x = a$.

A minimum (or local minimum) value of a function is the one which is less than all other values, on either side of the point in the immediate neighbourhood of the point.

Note : A neighbourhood of a point $x \in \mathbb{R}$ is defined by open interval $]x - \epsilon [,$ when $\epsilon > 0$.

25.5 CONDITIONS FOR MAXIMUM OR MINIMUM

We know that derivative of a function is positive when the function is increasing and the derivative is negative when the function is decreasing. We shall apply this result to find the condition for maximum or a function to have a minimum. Refer to Fig. 25.6, points B,D, F are points of maxima and points A,C,E are points of minima.

Now, on the left of B, the function is increasing and so $f'(x) > 0$, but on the right of B, the function is decreasing and, therefore, $f'(x) < 0$. This can be achieved only when $f'(x)$ becomes zero somewhere in between. We can rewrite this as follows :

A function $f(x)$ has a maximum value at a point if (i) $f'(x) = 0$ and (ii) $f'(x)$ changes sign from positive to negative in the neighbourhood of the point at which $f'(x) = 0$ (points taken from left to right).

Now, on the left of C (See Fig. 25.6), function is decreasing and $f'(x)$ therefore, is negative and on the right of C, $f(x)$ is increasing and so $f'(x)$ is positive. Once again $f'(x)$ will be zero before having positive values. We rewrite this as follows :

A function $f(x)$ has a minimum value at a point if (i) $f'(x) = 0$, and (ii) $f'(x)$ changes sign from negative to positive in the neighbourhood of the point at which $f'(x) = 0$.

Maxima and Minima

We should note here that $f'(x) = 0$ is necessary condition and is not a sufficient condition for maxima or minima to exist. We can have a function which is increasing, then constant and then again increasing function. In this case, $f'(x)$ does not change sign. The value for which $f'(x) = 0$ is not a point of maxima or minima. Such point is called point of inflexion.

For example, for the function $f(x) = x^3$, $x = 0$ is the point of inflexion as $f'(x) = 3x^2$ does not change sign as x passes through 0. $f'(x)$ is positive on both sides of the value '0' (tangents make acute angles with x -axis) (See Fig. 25.9).

Hence $f(x) = x^3$ has a point of inflexion at $x = 0$.

The points where $f'(x) = 0$ are called stationary points as the rate of change of the function is zero there. Thus points of maxima and minima are stationary points.




Fig. 25.9

MODULE - V Calculus



Notes

Remarks

The stationary points at which the function attains either local maximum or local minimum values are also called extreme points and both local maximum and local minimum values are called extreme values of $f(x)$. Thus a function attains an extreme value at $x=a$ if $f(a)$ is either a local maximum or a local minimum.

25.6 METHOD OF FINDING MAXIMA OR MINIMA

We have arrived at the method of finding the maxima or minima of a function. It is as follows :

- (i) Find $f'(x)$
- (ii) Put $f'(x) = 0$ and find stationary points
- (iii) Consider the sign of $f'(x)$ in the neighbourhood of stationary points. If it changes sign from +ve to -ve, then $f(x)$ has maximum value at that point and if $f'(x)$ changes sign from -ve to +ve, then $f(x)$ has minimum value at that point.
- (iv) If $f'(x)$ does not change sign in the neighbourhood of a point then it is a point of inflexion.

Example 25.7 Find the maximum (local maximum) and minimum (local minimum) points of the function $f(x) = x^3 - 3x^2 - 9x$.

Solution : Here $f(x) = x^3 - 3x^2 - 9x$

$$f'(x) = 3x^2 - 6x - 9$$

Step I. Now $f'(x) = 0$ gives us $3x^2 - 6x - 9 = 0$

or $x^2 - 2x - 3 = 0$

or $(x - 3)(x + 1) = 0$

or $x = 3, -1$

\therefore Stationary points are $x = 3, x = -1$

MODULE - V
Calculus

Notes

Step II. At

$x = 3$

For

$x < 3$

$f'(x) < 0$

and for

$x > 3$

$f'(x) > 0$

 $\therefore f'(x)$ changes sign from -ve to +ve in the neighbourhood of 3. $\therefore f(x)$ has minimum value at $x = 3$.**Step III.** At

$x = -1,$

For

$x < -1,$

$f'(x) > 0$

and for

$x > -1,$

$f'(x) < 0$

 $\therefore f'(x)$ changes sign from +ve to -ve in the neighbourhood of -1. $\therefore f(x)$ has maximum value at $x = -1$. $\therefore x = -1$ and $x = 3$ give us points of maxima and minima respectively. If we want to find maximum value (minimum value), then we have

$$\begin{aligned} \text{maximum value} &= f(-1) = (-1)^3 - 3(-1)^2 - 9(-1) \\ &= -1 - 3 + 9 = 5 \end{aligned}$$

and

$$\text{minimum value} = f(3) = 3^3 - 3(3)^2 - 9(3) = -27$$

 $\therefore (-1, 5)$ and $(3, -27)$ are points of local maxima and local minima respectively.**Example 25.8**

Find the local maximum and the local minimum of the function

$$f(x) = x^2 - 4x$$

Solution :

$$f(x) = x^2 - 4x$$

 \therefore

$$f'(x) = 2x - 4 = 2(x - 2)$$

Putting $f'(x) = 0$ yields $2x - 4 = 0$, i.e., $x = 2$.We have to examine whether $x = 2$ is the point of local maximum or local minimum or neither maximum nor minimum.Let us take $x = 1.9$ which is to the left of 2 and $x = 2.1$ which is to the right of 2 and find $f'(x)$ at these points.


$$f'(1.9) = 2(1.9 - 2) < 0$$

$$f'(2.1) = 2(2.1 - 2) > 0$$

Since $f'(x) < 0$ as we approach 2 from the left and $f'(x) > 0$ as we approach 2 from the right, therefore, there is a local minimum at $x = 2$.We can put our findings for sign of derivatives of $f(x)$ in any tabular form including the one given below :



Notes



Example 25.9 Find all local maxima and local minima of the function

$$f(x) = 2x^3 - 3x^2 - 12x + 8$$

Solution :

$$f(x) = 2x^3 - 3x^2 - 12x + 8$$

\therefore

$$f'(x) = 6x^2 - 6x - 12$$

$$= 6(x^2 - x - 2)$$

\therefore

$$f'(x) = 6(x + 1)(x - 2)$$

Now solving $f'(x) = 0$ for x , we get

$$6(x + 1)(x - 2) = 0$$

\Rightarrow

$$x = -1, 2$$

Thus

$$f'(x) = 0 \text{ at } x = -1, 2.$$

We examine whether these points are points of local maximum or local minimum or neither of them.

Consider the point $x = -1$

Let us take $x = -1.1$ which is to the left of -1 and $x = -0.9$ which is to the right of -1 and find $f'(x)$ at these points.

$$f'(-1.1) = 6(-1.1 + 1)(-1.1 - 2), \text{ which is positive i.e. } f'(x) > 0$$

$$f'(-0.9) = 6(-0.9 + 1)(-0.9 - 2), \text{ which is negative i.e. } f'(x) < 0$$

Thus, at $x = -1$, there is a local maximum.

Consider the point $x = 2$.

Now, let us take $x = 1.9$ which is to the left of $x = 2$ and $x = 2.1$ which is to the right of $x = 2$ and find $f'(x)$ at these points.

$$f'(1.9) = 6(1.9 + 1)(1.9 - 2)$$

$$= 6 \times (\text{Positive number}) \times (\text{negative number})$$


$$= \text{a negative number}$$

MODULE - V
Calculus



Notes

i.e. $f'(1.9) < 0$
and $f'(2.1) = 6(2.1+1)(2.1-2)$, which is positive
i.e., $f'(2.1) > 0$



$\therefore f'(x) < 0$ as we approach 2 from the left
and $f'(x) > 0$ as we approach 2 from the right.
 $\therefore x = 2$ is the point of local minimum

Thus $f(x)$ has local maximum at $x = -1$, maximum value of $f(x) = -2 - 3 + 12 + 8 = 15$
 $f(x)$ has local minimum at $x = 2$, minimum value of $f(x) = 2(8) - 3(4) - 12(2) + 8 = -12$

Sign of $f'(x)$

Point $x = -1$			Point $x = 2$	
Left of -1	Right of -1		Left of 2	Right of 2
positive	negative		negative	positive
local maximum			local minimum	

Example 25.10 Find local maximum and local minimum, if any, of the following function

$$f(x) = \frac{x}{1+x^2}$$

Solution :

$$f(x) = \frac{x}{1+x^2}$$

Then

$$f'(x) = \frac{(1+x^2)1 - (2x)x}{(1+x^2)^2}$$

$$= \frac{1-x^2}{(1+x^2)^2}$$

For finding points of local maximum or local minimum, equate $f'(x)$ to 0.

i.e.
$$\frac{1-x^2}{(1+x^2)^2} = 0$$



Notes

$$\Rightarrow 1 - x^2 = 0$$

$$\text{or } (1+x)(1-x) = 0 \quad \text{or} \quad x = 1, -1.$$

Consider the value $x = 1$.

The sign of $f'(x)$ for values of x slightly less than 1 and slightly greater than 1 changes from positive to negative. Therefore there is a local maximum at $x = 1$, and the local maximum

$$\text{value} = \frac{1}{1+(1)^2} = \frac{1}{1+1} = \frac{1}{2}$$

Now consider $x = -1$.

$f'(x)$ changes sign from negative to positive as x passes through -1 , therefore, $f(x)$ has a local minimum at $x = -1$

$$\text{Thus, the local minimum value} = \frac{-1}{2}$$

Example 25.11 Find the local maximum and local minimum, if any, for the function

$$f(x) = \sin x + \cos x, 0 < x < \frac{\pi}{2}$$

Solution : We have $f(x) = \sin x + \cos x$

$$f'(x) = \cos x - \sin x$$

For local maxima/minima, $f'(x) = 0$

$$\therefore \cos x - \sin x = 0$$

$$\text{or, } \tan x = 1 \quad \text{or, } x = \frac{\pi}{4} \text{ in } 0 < x < \frac{\pi}{2}$$

$$\text{At } x = \frac{\pi}{4},$$

$$\text{For } x < \frac{\pi}{4}, \cos x > \sin x$$

$$\therefore f'(x) = \cos x - \sin x > 0$$

$$\text{For } x > \frac{\pi}{4}, \cos x - \sin x < 0$$

$$\therefore f'(x) = \cos x - \sin x < 0$$

$\therefore f'(x)$ changes sign from positive to negative in the neighbourhood of $\frac{\pi}{4}$.

$\therefore x = \frac{\pi}{4}$ is a point of local maxima.

$$\text{Maximum value} = f\left(\frac{\pi}{4}\right) = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

\therefore Point of local maxima is $\left(\frac{\pi}{4}, \sqrt{2}\right)$

MODULE - V
Calculus**CHECK YOUR PROGRESS 25.3**

Notes

Find all points of local maxima and local minima of the following functions. Also, find the maxima and minima at such points.

- | | |
|------------------------------|--------------------------------|
| 1. $x^2 - 8x + 12$ | 2. $x^3 - 6x^2 + 9x + 15$ |
| 3. $2x^3 - 21x^2 + 36x - 20$ | 4. $x^4 - 62x^2 + 120x + 9$ |
| 5. $(x - 1)(x - 2)^2$ | 6. $\frac{x - 1}{x^2 + x + 2}$ |

25.7 USE OF SECOND DERIVATIVE FOR DETERMINATION OF MAXIMUM AND MINIMUM VALUES OF A FUNCTION

We now give below another method of finding local maximum or minimum of a function whose second derivative exists. Various steps used are :

- Let the given function be denoted by $f(x)$.
- Find $f'(x)$ and equate it to zero.
- Solve $f'(x) = 0$, let one of its real roots be $x = a$.
- Find its second derivative, $f''(x)$. For every real value 'a' of x obtained in step (iii), evaluate $f''(a)$. Then if

$f''(a) < 0$ then $x = a$ is a point of local maximum.

$f''(a) > 0$ then $x = a$ is a point of local minimum.

$f''(a) = 0$ then we use the sign of $f'(x)$ on the left of 'a' and on the right of 'a' to arrive at the result.

Example 25.12 Find the local minimum of the following function :

$$2x^3 - 21x^2 + 36x - 20$$

Solution : Let $f(x) = 2x^3 - 21x^2 + 36x - 20$

Then $f'(x) = 6x^2 - 42x + 36$

$$= 6(x^2 - 7x + 6)$$

$$= 6(x - 1)(x - 6)$$

For local maximum or minimum

$$f'(x) = 0$$

or $6(x - 1)(x - 6) = 0 \Rightarrow x = 1, 6$



Notes

$$\begin{aligned} \text{Now} \quad f''(x) &= \frac{d}{dx} f'(x) \\ &= \frac{d}{dx} [6(x^2 - 7x + 6)] \\ &= 12x - 42 \\ &= 6(2x - 7) \end{aligned}$$

For $x = 1$, $f''(1) = 6(2 \cdot 1 - 7) = -30 < 0$
 $x = 1$ is a point of local maximum.

and $f(1) = 2(1)^3 - 21(1)^2 + 36(1) - 20 = 3$ is a local maximum.

For $x = 6$,

$$f''(6) = 6(2 \cdot 6 - 7) = 30 > 0$$

$\therefore x = 6$ is a point of local minimum

and $f(6) = 2(6)^3 - 21(6)^2 + 36(6) - 20 = 128$ is a local minimum.

Example 25.13 Find local maxima and minima (if any) for the function

$$f(x) = \cos 4x; \quad 0 < x < \frac{\pi}{2}$$

Solution :

$$f(x) = \cos 4x$$

$$\therefore f'(x) = -4 \sin 4x$$

$$\text{Now, } f'(x) = 0 \quad \Rightarrow \quad -4 \sin 4x = 0$$

$$\text{or, } \sin 4x = 0 \quad \text{or, } \quad 4x = 0, \pi, 2\pi$$

$$\text{or, } \quad x = 0, \frac{\pi}{4}, \frac{\pi}{2}$$

$$\therefore \quad x = \frac{\pi}{4} \quad \left[\because 0 < x < \frac{\pi}{2} \right]$$

$$\text{Now, } f''(x) = -16 \cos 4x$$

$$\begin{aligned} \text{at } x = \frac{\pi}{4}, f''(x) &= -16 \cos \pi \\ &= -16(-1) = 16 > 0 \end{aligned}$$

$\therefore f(x)$ is minimum at $x = \frac{\pi}{4}$

$$\text{Minimum value } f\left(\frac{\pi}{4}\right) = \cos \pi = -1$$

MODULE - V
Calculus

Notes

Example 25.14 (a) Find the maximum value of $2x^3 - 24x + 107$ in the interval $[-3, -1]$.
(b) Find the minimum value of the above function in the interval $[1, 3]$.

Solution : Let $f(x) = 2x^3 - 24x + 107$

$$f'(x) = 6x^2 - 24$$

For local maximum or minimum,

$$f'(x) = 0$$

$$\text{i.e. } 6x^2 - 24 = 0 \quad \Rightarrow \quad x = -2, 2$$

Out of two points obtained on solving $f'(x) = 0$, only -2 belong to the interval $[-3, -1]$. We shall, therefore, find maximum if any at $x = -2$ only.

$$\text{Now } f''(x) = 12x$$

$$\therefore f''(-2) = 12(-2) = -24$$

$$\text{or } f''(-2) < 0$$

which implies the function $f(x)$ has a maximum at $x = -2$.

$$\begin{aligned} \therefore \text{ Required maximum value} &= 2(-2)^3 - 24(-2) + 107 \\ &= 139 \end{aligned}$$

Thus the point of maximum belonging to the given interval $[-3, -1]$ is -2 and, the maximum value of the function is 139.

$$\text{(b) Now } f''(x) = 12x$$

$$\therefore f''(2) = 24 > 0, \quad [\because 2 \text{ lies in } [1, 3]]$$

which implies, the function $f(x)$ shall have a minimum at $x = 2$.

$$\begin{aligned} \therefore \text{ Required minimum} &= 2(2)^3 - 24(2) + 107 \\ &= 75 \end{aligned}$$

Example 25.15 Find the maximum and minimum value of the function

$$f(x) = \sin x (1 + \cos x) \text{ in } (0, \pi).$$

Solution : We have, $f(x) = \sin x (1 + \cos x)$

$$f'(x) = \cos x (1 + \cos x) + \sin x (-\sin x)$$

$$= \cos x + \cos^2 x - \sin^2 x$$

$$= \cos x + \cos^2 x - (1 - \cos^2 x) = 2\cos^2 x + \cos x - 1$$

For stationary points, $f'(x) = 0$

$$\therefore 2\cos^2 x + \cos x - 1 = 0$$

$$\therefore \cos x = \frac{-1 \pm \sqrt{1+8}}{4} = \frac{-1 \pm 3}{4} = -1, \frac{1}{2}$$



Notes

$$\therefore x = \pi, \frac{\pi}{3}$$

Now, $f(0) = 0$

$$f\left(\frac{\pi}{3}\right) = \sin \frac{\pi}{3} \left(1 + \cos \frac{\pi}{3}\right) = \frac{\sqrt{3}}{2} \left(1 + \frac{1}{2}\right) = \frac{3\sqrt{3}}{4}$$

and $f(\pi) = 0$

$$\therefore f(x) \text{ has maximum value } \frac{3\sqrt{3}}{4} \text{ at } x = \frac{\pi}{3}$$

and minimum value 0 at $x = 0$ and $x = \pi$.



CHECK YOUR PROGRESS 25.4

Find local maximum and local minimum for each of the following functions using second order derivatives.

1. $2x^3 + 3x^2 - 36x + 10$

2. $-x^3 + 12x^2 - 5$

3. $(x-1)(x+2)^2$

4. $x^5 - 5x^4 + 5x^3 - 1$

5. $\sin x (1 + \cos x), 0 < x < \frac{\pi}{2}$

6. $\sin x + \cos x, 0 < x < \frac{\pi}{2}$

7. $\sin 2x - x, \frac{-\pi}{2} \leq x \leq \frac{\pi}{2}$

25.8 APPLICATIONS OF MAXIMA AND MINIMA TO PRACTICAL PROBLEMS

The application of derivative is a powerful tool for solving problems that call for minimising or maximising a function. In order to solve such problems, we follow the steps in the following order :

- (i) Frame the function in terms of variables discussed in the data.
- (ii) With the help of the given conditions, express the function in terms of a single variable.
- (iii) Lastly, apply conditions of maxima or minima as discussed earlier.

Example 25.16 Find two positive real numbers whose sum is 70 and their product is maximum.

Solution : Let one number be x . As their sum is 70, the other number is $70-x$. As the two numbers are positive, we have, $x > 0, 70-x > 0$

$$70 - x > 0 \quad \Rightarrow \quad x < 70$$

$$\therefore 0 < x < 70$$

MODULE - V
Calculus


Notes

Let their product be $f(x)$

Then
$$f(x) = x(70 - x) = 70x - x^2$$

We have to maximize the product $f(x)$.

We, therefore, find $f'(x)$ and put that equal to zero.

$$f'(x) = 70 - 2x$$

For maximum product, $f'(x) = 0$

or
$$70 - 2x = 0$$

or
$$x = 35$$

Now $f''(x) = -2$ which is negative. Hence $f(x)$ is maximum at $x = 35$

The other number is $70 - x = 35$

Hence the required numbers are 35, 35.

Example 25.17 Show that among rectangles of given area, the square has the least perimeter.

Solution : Let x, y be the length and breadth of the rectangle respectively.

\therefore Its area = xy

Since its area is given, represent it by A , so that we have

$$A = xy$$

or
$$y = \frac{A}{x} \quad \dots (i)$$

Now, perimeter say P of the rectangle = $2(x + y)$

or
$$P = 2\left(x + \frac{A}{x}\right)$$

\therefore
$$\frac{dP}{dx} = 2\left(1 - \frac{A}{x^2}\right) \quad \dots(ii)$$

For a minimum P , $\frac{dP}{dx} = 0$.

i.e.
$$2\left(1 - \frac{A}{x^2}\right) = 0$$

or
$$A = x^2 \quad \text{or} \quad \sqrt{A} = x$$

Now,
$$\frac{d^2P}{dx^2} = \frac{4A}{x^3}$$
, which is positive.

Hence perimeter is minimum when $x = \sqrt{A}$

\therefore
$$y = \frac{A}{x}$$



Notes

$$= \frac{x^2}{x} = x \quad (\because A = x^2)$$

Thus, the perimeter is minimum when rectangle is a square.

Example 25.18 An open box with a square base is to be made out of a given quantity of sheet of area a^2 . Show that the maximum volume of the box is $\frac{a^3}{6\sqrt{3}}$.

Solution : Let x be the side of the square base of the box and y its height.

Total surface area of othe box $= x^2 + 4xy$

$$\therefore x^2 + 4xy = a^2 \quad \text{or} \quad y = \frac{a^2 - x^2}{4x}$$

Volume of the box, $V = \text{base area} \times \text{height}$

$$= x^2 y = x^2 \left(\frac{a^2 - x^2}{4x} \right)$$

$$\text{or} \quad V = \frac{1}{4} (a^2 x - x^3) \quad \dots(i)$$

$$\therefore \frac{dV}{dx} = \frac{1}{4} (a^2 - 3x^2)$$

For maxima/minima $\frac{dV}{dx} = 0$

$$\therefore \frac{1}{4} (a^2 - 3x^2) = 0$$

$$x^2 = \frac{a^2}{3} \Rightarrow x = \frac{a}{\sqrt{3}} \quad \dots(ii)$$

From (i) and (ii), we get

$$\text{Volume} = \frac{1}{4} \left(\frac{a^3}{\sqrt{3}} - \frac{a^3}{3\sqrt{3}} \right) = \frac{a^3}{6\sqrt{3}} \quad \dots(iii)$$

$$\text{Again} \quad \frac{d^2V}{dx^2} = \frac{d}{dx} \frac{1}{4} (a^2 - 3x^2) = -\frac{3}{2} x$$

x being the length of the side, is positive.

$$\therefore \frac{d^2V}{dx^2} < 0$$

\therefore The volume is maximum.

Hence maximum volume of the box $= \frac{a^3}{6\sqrt{3}}$.

MODULE - V
Calculus



Notes

Example 25.19 Show that of all rectangles inscribed in a given circle, the square has the maximum area.

Solution : Let ABCD be a rectangle inscribed in a circle of radius r. Then diameter AC = 2r

Let AB = x and BC = y

Then $AB^2 + BC^2 = AC^2$ or $x^2 + y^2 = (2r)^2 = 4r^2$

Now area A of the rectangle = xy

$\therefore A = x\sqrt{4r^2 - x^2}$

$\therefore \frac{dA}{dx} = \frac{x(-2x)}{2\sqrt{4r^2 - x^2}} + \sqrt{4r^2 - x^2}$
 $= \frac{4r^2 - 2x^2}{\sqrt{4r^2 - x^2}}$

For maxima/minima, $\frac{dA}{dx} = 0$

$\frac{4r^2 - 2x^2}{\sqrt{4r^2 - x^2}} = 0 \Rightarrow x = \sqrt{2}r$

Now

$\frac{d^2A}{dx^2} = \frac{\sqrt{4r^2 - x^2}(-4x) - (4r^2 - 2x^2)\frac{(-2x)}{2\sqrt{4r^2 - x^2}}}{(4r^2 - x^2)}$
 $= \frac{-4x(4r^2 - x^2) + x(4r^2 - 2x^2)}{(4r^2 - x^2)^{\frac{3}{2}}}$
 $= \frac{-4\sqrt{2}(2r^2) + 0}{(2r^2)^{\frac{3}{2}}}$
 $= \frac{-8\sqrt{2}r^3}{2\sqrt{2}r^3} = -4 < 0$




Fig. 25.10

... (Putting $x = \sqrt{2}r$)

Thus, A is maximum when $x = \sqrt{2}r$

Now, from (i), $y = \sqrt{4r^2 - 2r^2} = \sqrt{2}r$

$x = y$. Hence, rectangle ABCD is a square.



Notes

Example 25.20 Show that the height of a closed right circular cylinder of a given volume and least surface is equal to its diameter.

Solution : Let V be the volume, r the radius and h the height of the cylinder.

Then,
$$V = \pi r^2 h$$

or
$$h = \frac{V}{\pi r^2} \quad \dots (i)$$

Now surface area
$$S = 2\pi r h + 2\pi r^2$$

$$= 2\pi r \cdot \frac{V}{\pi r^2} + 2\pi r^2 = \frac{2V}{r} + 2\pi r^2$$

Now
$$\frac{dS}{dr} = \frac{-2V}{r^2} + 4\pi$$

For minimum surface area, $\frac{dS}{dr} = 0$

$\therefore \frac{-2V}{r^2} + 4\pi = 0$

or
$$V = 2\pi r^3$$

From (i) and (ii), we get
$$h = \frac{2\pi r^3}{\pi r^2} = 2r \quad \dots(ii)$$

Again,
$$\frac{d^2S}{dr^2} = \frac{4V}{r^3} + 4\pi = \frac{4 \cdot 2\pi r^3}{r^3} + 4\pi$$

$$= 12\pi > 0$$

... [Using (ii)]

$\therefore S$ is least when $h = 2r$

Thus, height of the cylinder = diameter of the cylinder.

Example 25.21 Show that a closed right circular cylinder of given surface has maximum volume if its height equals the diameter of its base.

Solution : Let S and V denote the surface area and the volume of the closed right circular cylinder of height h and base radius r .

Then
$$S = 2\pi r h + 2\pi r^2 \quad \dots(i)$$

(Here surface is a constant quantity, being given)

$$V = \pi r^2 h$$

$\therefore V = \pi r^2 \left[\frac{S - 2\pi r^2}{2\pi r} \right]$

$$= \frac{r}{2} [S - 2\pi r^2]$$

MODULE - V
Calculus



Notes

$$V = \frac{Sr}{2} - \pi r^3$$

$$\frac{dV}{dr} = \frac{S}{2} - \pi(3r^2)$$

For maximum or minimum, $\frac{dV}{dr} = 0$

i.e.,
$$\frac{S}{2} - \pi(3r^2) = 0$$

or
$$S = 6\pi r^2$$

From (i), we have
$$6\pi r^2 = 2\pi rh + 2\pi r^2$$

$\Rightarrow 4\pi r^2 = 2\pi rh$

$\Rightarrow 2r = h \quad \dots(ii)$

Also,
$$\frac{d^2V}{dr^2} = \frac{d}{dr} \left[\frac{S}{2} - 3\pi r^2 \right]$$

$$= -6\pi,$$

= a negative quantity

$$\therefore \frac{d}{dr} \left(\frac{S}{2} \right) = 0$$

Hence the volume of the right circular cylinder is maximum when its height is equal to twice its radius i.e. when $h = 2r$.




Fig. 25.11

Example 25.22 A square metal sheet of side 48 cm. has four equal squares removed from the corners and the sides are then turned up so as to form an open box. Determine the size of the square cut so that volume of the box is maximum.

Solution : Let the side of each of the small squares cut be x cm, so that each side of the box to be made is $(48 - 2x)$ cm. and height x cm.

Now $x > 0$, $48 - 2x > 0$, i.e. $x < 24$

$\therefore x$ lies between 0 and 24 or $0 < x < 24$

Now, Volume V of the box

$$= (48 - 2x)(48 - 2x)x$$

i.e.
$$V = (48 - 2x)^2 \cdot x$$

$\therefore \frac{dV}{dx} = (48 - 2x)^2 + 2(48 - 2x)(-2)x$

$$= (48 - 2x)(48 - 6x)$$




Fig. 25.12



Notes

Condition for maximum or minimum is $\frac{dV}{dx} = 0$

i.e., $(48 - 2x)(48 - 6x) = 0$

We have either $x = 24$, or $x = 8$

$\therefore 0 < x < 24$

\therefore Rejecting $x = 24$, we have, $x = 8$ cm.

Now, $\frac{d^2V}{dx^2} = 24x - 384$

$$\left(\frac{d^2V}{dx^2}\right)_{x=8} = 192 - 384 = -192 < 0,$$

Hence for $x = 8$, the volume is maximum.

Hence the square of side 8 cm. should be cut from each corner.

Example 25.23 The profit function $P(x)$ of a firm, selling x items per day is given by

$$P(x) = (150 - x)x - 1625.$$

Find the number of items the firm should manufacture to get maximum profit. Find the maximum profit.

Solution : It is given that 'x' is the number of items produced and sold out by the firm every day. In order to maximize profit,

$$P'(x) = 0 \text{ i.e. } \frac{dP}{dx} = 0$$

or $\frac{d}{dx} [(150 - x)x - 1625] = 0$

or $150 - 2x = 0$

or $x = 75$

Now, $\frac{d}{dx} P'(x) = P''(x) = -2$ a negative quantity. Hence $P(x)$ is maximum for $x = 75$.

Thus, the firm should manufacture only 75 items a day to make maximum profit.

$$\begin{aligned} \text{Now, Maximum Profit} &= P(75) = (150 - 75)75 - 1625 \\ &= \text{Rs. } (75 \times 75 - 1625) \\ &= \text{Rs. } (5625 - 1625) \\ &= \text{Rs. } 4000 \end{aligned}$$

MODULE - V
Calculus



Notes

Example 25.24 Find the volume of the largest cylinder that can be inscribed in a sphere of radius 'r' cm.

Solution : Let h be the height and R the radius of the base of the inscribed cylinder. Let V be the volume of the cylinder.

Then $V = \pi R^2 h$... (i)

From ΔOCB , we have

$$r^2 = \left(\frac{h}{2}\right)^2 + R^2 \quad \dots (\because OB^2 = OC^2 + BC^2)$$

$\therefore R^2 = r^2 - \frac{h^2}{4}$... (ii)

Now $V = \pi \left(r^2 - \frac{h^2}{4} \right) h = \pi^2 h - \frac{\pi h^3}{4}$

$\therefore \frac{dV}{dh} = \pi^2 - \frac{3\pi h^2}{4}$

For maxima/minima, $\frac{dV}{dh} = 0$

$\therefore \pi r^2 - \frac{3\pi h^2}{4} = 0$

$\Rightarrow h^2 = \frac{4r^2}{3} \Rightarrow h = \frac{2r}{\sqrt{3}}$

Now $\frac{d^2V}{dh^2} = -\frac{3\pi h}{2}$

$\therefore \frac{d^2V}{dh^2} \left(\text{at } h = \frac{2r}{\sqrt{3}} \right) = \frac{3\pi \times 2r}{2 \times \sqrt{3}} = -\sqrt{3} \pi < 0$

$\therefore V$ is maximum at $h = \frac{2r}{\sqrt{3}}$

Putting $h = \frac{2r}{\sqrt{3}}$ in (ii), we get

$$R^2 = r^2 - \frac{4r^2}{4 \times 3} = \frac{2r^2}{3}$$

$\therefore R = \sqrt{\frac{2}{3}} r$

Maximum volume of the cylinder = $\pi R^2 h$

$$= \pi \left(\frac{2}{3} r^2 \right) \frac{2r}{\sqrt{3}} = \frac{4\pi r^3}{3\sqrt{3}} \text{ cm}^3.$$




Fig. 25.13

**CHECK YOUR PROGRESS 25.5**

Notes

- Find two numbers whose sum is 15 and the square of one multiplied by the cube of the other is maximum.
- Divide 15 into two parts such that the sum of their squares is minimum.
- Show that among the rectangles of given perimeter, the square has the greatest area.
- Prove that the perimeter of a right angled triangle of given hypotenuse is maximum when the triangle is isosceles.
- A window is in the form of a rectangle surmounted by a semi-circle. If the perimeter be 30 m, find the dimensions so that the greatest possible amount of light may be admitted.
- Find the radius of a closed right circular cylinder of volume 100 c.c. which has the minimum total surface area.
- A right circular cylinder is to be made so that the sum of its radius and its height is 6 m. Find the maximum volume of the cylinder.
- Show that the height of a right circular cylinder of greatest volume that can be inscribed in a right circular cone is one-third that of the cone.
- A conical tent of the given capacity (volume) has to be constructed. Find the ratio of the height to the radius of the base so as to minimise the canvas required for the tent.
- A manufacturer needs a container that is right circular cylinder with a volume 16π cubic meters. Determine the dimensions of the container that uses the least amount of surface (sheet) material.
- A movie theatre's management is considering reducing the price of tickets from Rs.55 in order to get more customers. After checking out various facts they decide that the average number of customers per day 'q' is given by the function where x is the amount of ticket price reduced. Find the ticket price that result in maximum revenue.

$$q = 500 + 100x$$

where x is the amount of ticket price reduced. Find the ticket price that result is maximum revenue.

**LET US SUM UP**

- Increasing function :** A function $f(x)$ is said to be increasing in the closed interval $[a,b]$ if $f(x_2) \geq f(x_1)$ whenever $x_2 > x_1$
- Decreasing function :** A function $f(x)$ is said to be decreasing in the closed interval $[a,b]$ if $f(x_2) \leq f(x_1)$ whenever $x_2 > x_1$
- $f(x)$ is increasing in an open interval $]a,b[$ if $f'(x) > 0$ for all $x \in]a,b[$
- $f(x)$ is decreasing in an open interval $]a,b[$

MODULE - V
Calculus

Notes

if $f'(x) < 0$ for all $x \in [a, b]$

- **Monotonic function :**

- (i) A function is said to be monotonic (increasing) if it increases in the given interval.
- (ii) A function $f(x)$ is said to be monotonic (decreasing) if it decreases in the given interval.

A function $f(x)$ which increases and decreases in a given interval, is not monotonic.

- In an interval around the point $x = a$ of the function $f(x)$,
 - (i) if $f'(x) > 0$ on the left of the point 'a' and $f'(x) < 0$ on the right of the point $x = a$, then $f(x)$ has a local maximum.
 - (ii) if $f'(x) < 0$ on the left of the point 'a' and $f'(x) > 0$ on the right of the point $x = a$, then $f(x)$ has a local minimum.
- If $f(x)$ has a local maximum or local minimum at $x = a$ and $f(x)$ is derivable at $x = a$, then $f'(a) = 0$
- (i) If $f'(x)$ changes sign from positive to negative as x passes through 'a', then $f(x)$ has a local maximum at $x = a$.
- (ii) If $f'(x)$ changes sign from negative to positive as x passes through 'a', then $f(x)$ has a local minimum at $x = a$.
- **Second order derivative Test :**
 - (i) If $f'(a) = 0$, and $f''(a) < 0$; then $f(x)$ has a local maximum at $x = a$.
 - (ii) If $f'(a) = 0$, and $f''(a) > 0$; then $f(x)$ has a local minimum at $x = a$.
 - (iii) In case $f'(a) = 0$, and $f''(a) = 0$; then to determine maximum or minimum at $x = a$, we use the method of change of sign of $f'(x)$ as x passes through 'a' to.

**SUPPORTIVE WEB SITES**

- <http://www.wikipedia.org>
- <http://mathworld.wolfram.com>

**TERMINAL EXERCISE**

1. Show that $f(x) = x^2$ is neither increasing nor decreasing for all $x \in \mathbb{R}$.
Find the intervals for which the following functions are increasing or decreasing :
2. $2x^3 - 3x^2 - 12x + 6$
3. $\frac{x}{4} + \frac{4}{x}, x \neq 0$
4. $x^4 - 2x^2$
5. $\sin x - \cos x, 0 \leq x \leq 2\pi$

Maxima and Minima

Find the local maxima or minima of the following functions :

6. (a) $x^3 - 6x^2 + 9x + 7$ (b) $2x^3 - 24x + 107$
(c) $x^3 + 4x^2 - 3x + 2$ (d) $x^4 - 62x^2 + 120x + 9$
7. (a) $\frac{1}{x^2 + 2}$ (b) $\frac{x}{(x-1)(x-4)}, 1 < x < 4$
(c) $x\sqrt{1-x}, x < 1$
8. (a) $\sin x + \frac{1}{2}\cos 2x, 0 \leq x \leq \frac{\pi}{2}$ (b) $\sin 2x, 0 \leq x \leq 2\pi$
(c) $-x + 2\sin x, 0 \leq x \leq 2\pi$
9. For what value of x lying in the closed interval $[0,5]$, the slope of the tangent to
$$x^3 - 6x^2 + 9x + 4$$
is maximum. Also, find the point.
10. Find the value of the greatest slope of a tangent to
$$-x^3 + 3x^2 + 2x - 27$$
 at a point of the curve. Find also the point.
11. A container is to be made in the shape of a right circular cylinder with total surface area of 24π sq. m. Determine the dimensions of the container if the volume is to be as large as possible.
12. A hotel complex consisting of 400 two bedroom apartments has 300 of them rented and the rent is Rs. 360 per day. Management's research indicates that if the rent is reduced by x rupees then the number of apartments rented q will be $q = \frac{5}{4}x + 300, 0 \leq x \leq 80$. Determine the rent that results in maximum revenue. Also find the maximum revenue.

MODULE - V Calculus



Notes

**CHECK YOUR PROGRESS 25.2**

1. (a) Increasing for $x > \frac{7}{2}$, Decreasing for $x < \frac{7}{2}$
- (b) Increasing for $x > \frac{5}{2}$, Decreasing for $x < \frac{5}{2}$
2. (a) Increasing for $x > 6$ or $x < -2$, Decreasing for $-2 < x < 6$
- (b) Increasing for $x > 4$ or $x < 2$, Decreasing for x in the interval $]2,4[$
3. (a) Increasing for $x < -2$; decreasing for $x > -2$
- (b) Increasing in the interval $-1 < x < 2$, Decreasing for $x > -1$ or $x < -2$
4. (a) Increasing always.
- (b) Increasing for $x > 2$, Decreasing in the interval $0 < x < 2$
- (c) Increasing for $x > 2$ or $x < -2$ Decreasing in the interval $-2 < x < 2$
5. (c) Increasing in the interval $\frac{3\pi}{8} \leq x \leq \frac{7\pi}{8}$
Decreasing in the interval $0 \leq x \leq \frac{3\pi}{8}$

Points at which the tangents are parallel to x-axis are $x = \frac{3\pi}{8}$ and $x = \frac{7\pi}{8}$

CHECK YOUR PROGRESS 25.3

1. Local minimum is -4 at $x = 4$
2. Local minimum is 15 at $x = 3$, Local maximum is 19 at $x = 1$.
3. Local minimum is -128 at $x = 6$, Local maximum is -3 at $x = 1$.
4. Local minimum is -1647 at $x = -6$, Local minimum is -316 at $x = 5$,
Local maximum is 68 at $x = 1$.
5. Local minimum at $x = 0$ is -4 , Local maximum at $x = -2$ is 0 .
6. Local minimum at $x = -1$, value = -1 Local maximum at $x = 3$, value = $\frac{1}{7}$

CHECK YOUR PROGRESS 25.4

1. Local minimum is -34 at $x = 2$, Local maximum is 91 at $x = -3$.
2. Local minimum is -5 at $x = 0$, Local maximum is 251 at $x = 8$.
3. Local minimum -4 at $x = 0$, Local maximum 0 at $x = -2$.
4. Local minimum = -28 ; $x = 3$, Local maximum = 0 ; $x = 1$.
Neither maximum nor minimum at $x = 0$.



Notes

5. Local maximum = $\frac{3\sqrt{3}}{4}$; $x = \frac{\pi}{3}$

6. Local maximum = $\sqrt{2}$; $x = \frac{\pi}{4}$

7. Local minimum = $-\frac{\sqrt{3}}{2} + \frac{\pi}{6}$; $x = \frac{\pi}{6}$, Local maximum = $\frac{\sqrt{3}}{2} - \frac{\pi}{6}$; $x = \frac{\pi}{6}$

CHECK YOUR PROGRESS 25.5

1. Numbers are 6,9.

2. Parts are 7.5, 7.5

5. Dimensions are : = $\frac{30}{\pi+4}$, $\frac{30}{\pi+4}$ meters each.

6. radius = $\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ cm; height = $2\left(\frac{50}{\pi}\right)^{\frac{1}{3}}$ cm

7. Maximum Volume = 32π cubic meters. 9. $h = \sqrt{2}r$

10. $r = 2$ meters, $h = 4$ meters. 11. Rs. 30,00

TERMINAL EXERCISE

2. Increasing for $x > 2$ or $x < -1$,

Decreasing in the interval $-1 < x < 2$

3. Increasing for $x > 4$ or $x < -4$,

Decreasing in the interval $]-4,4[$

4. Increasing for $x > 1$ or $-1 < x < 0$,

Decreasing for $x < -1$ or $0 < x < 1$

5. Increasing for $0 \leq x \leq \frac{3\pi}{4}$ or $\frac{7\pi}{4} \leq x \leq 2\pi$, Decreasing for $\frac{3\pi}{4} \leq x \leq \frac{7\pi}{4}$.

6. (a) Local maximum is 11 at $x = 1$; local minimum is 7 at $x = 3$.

(b) Local maximum is 139 at $x = -2$; local minimum is 75 at $x = 2$.

(c) Local maximum is 20 at $x = -3$; local minimum is $\frac{40}{27}$ at $x = \frac{1}{3}$.

(d) Local maximum is 68 at $x = 1$; local minimum is -316 at $x = 5$ and -1647 at $x = -6$.

7. (a) Local minimum is $\frac{1}{2}$ at $x = 0$. (b) Local maximum is -1 at $x = 2$.

(c) Local maximum is $\frac{2}{3\sqrt{3}}$ at $x = \frac{2}{3}$.

8. (a) Local maximum is $\frac{3}{4}$ at $x = \frac{\pi}{6}$; Local minimum is $\frac{1}{2}$ at $x = \frac{\pi}{2}$;

MODULE - V
Calculus


Notes

- (b) Local maximum is 1 at $x = \frac{\pi}{4}$ and $x = \frac{5\pi}{4}$; Local minimum is -1 at $x = \frac{3\pi}{4}$
- (c) Local maximum is $\frac{-\pi}{4} + \sqrt{3}$ at $x = \frac{\pi}{3}$; Local minimum is $\frac{-5\pi}{3} - \sqrt{3}$ at $x = \frac{5\pi}{3}$.
9. Greatest slope is 24 at $x = 5$; Coordinates of the point: (5, 24).
10. Greatest slope of a tangent is 5 at $x = 1$, The point is (1, -23)
11. Radius of base = 2 m, Height of cylinder = 4 m.
12. Rent reduced to Rs. 300, The maximum revenue = Rs. 1,12,500.